



Review Article

A prospective review of the health-promoting potential of Jing Si Herbal Tea

Tsung-Jung Ho^{a,b,c†}, Tanvir Ahmed^{d†}, Marthandam Asokan Shibu^{d,e}, Yu-Jung Lin^{b,d}, Cheng Yen Shih^{f,g}, Pi-Yu Lin^f, Shinn-Zong Ling^{g,h}, Chien-Yi Chiang^b, Wei-Wen Kuo^{i,j}, Chih-Yang Huang^{d,k,l,m,n*}

^aDepartment of Chinese Medicine, Hualien Tzu Chi Hospital, Buddhist Tzu Chi Medical Foundation and Tzu Chi University, Hualien, Taiwan, ^bSchool of Post-Baccalaureate Chinese Medicine, College of Medicine, Tzu Chi University, Hualien, Taiwan, ^cIntegration Center of Traditional Chinese and Modern Medicine, Hualien Tzu Chi Hospital, Buddhist Tzu Chi Medical Foundation, Hualien, Taiwan, ^dCardiovascular and Mitochondrial Related Disease Research Center, Hualien Tzu Chi Hospital, Buddhist Tzu Chi Medical Foundation, Hualien, Taiwan, ^eDepartment of Biotechnology, Bharathiar University, Coimbatore, Tamil Nadu, India, ^fBuddhist Compassion Relief Tzu Chi Foundation, Hualien, Taiwan, ^gBioinnovation Center, Buddhist Tzu Chi Medical Foundation, Hualien, Taiwan, ^hDepartment of Neurosurgery, Hualien Tzu Chi Hospital, Buddhist Tzu Chi Medical Foundation, Hualien, Taiwan, ⁱDepartment of Biological Science and Technology, College of Life Sciences, China Medical University, Taichung, Taiwan, ^jPh. D. Program for Biotechnology Industry, China Medical University, Taichung, Taiwan, ^kGraduate Institute of Biomedical Sciences, China Medical University, Taichung, Taiwan, ^lDepartment of Biological Science and Technology, Asia University, Taichung, Taiwan, ^mDepartment of Medical Research, China Medical University Hospital and China Medical University, Taichung, Taiwan, ⁿCenter of General Education, Buddhist Tzu Chi Medical Foundation, Tzu Chi University of Science and Technology, Hualien, Taiwan

† Both authors contributed equally to this work.

Submission : 09-Aug-2023
Revision : 29-Aug-2023
Acceptance : 02-Oct-2023
Web Publication : 12-Jan-2024

ABSTRACT

Traditional Chinese medicine (TCM) has gained considerable attention over the past few years for its multicomponent, multitarget, and multi-pathway approach to treating different diseases. Studies have shown that TCMs as adjuvant therapy along with conventional treatment may benefit in safely treating various disorders. However, investigations on finding effective herbal combinations are ongoing. A novel TCM formula, “Jing Si Herbal Tea (JSHT),” has been reported recently for their health-promoting effects in improving overall body and mental health. JSHT is a combination of eight herbs recognized in Chinese herbal pharmacopoeia for their anti-viral, anti-aging, and anti-cancer properties as well as protective effects against cardiovascular, metabolic, neural, digestive, and genitourinary diseases. Thus, to better understand the beneficial effects of the ingredients of JSHT on health, this review intends to summarize the preclinical and clinical studies of the ingredients of JSHT on human health and diseases, and possible therapeutic effects with the related mode of actions and future prospects for their application in complementary therapies.

KEYWORDS: Health benefits, Jing Si herbal tea, Pharmacological properties, Phytochemical compounds, Traditional Chinese medicine

INTRODUCTION

Recently, there has been a noticeable increase in the prevalence of complementary and alternative medicine (CAM) practice, and 75% of the world's population relies on herbs for their primary health-care requirements [1].

Access this article online

Quick Response Code:



Website: www.tcmjmed.com

DOI: 10.4103/tcmj.tcmj_194_23

***Address for correspondence:** Prof. Chih-Yang Huang, Cardiovascular and Mitochondria Related Diseases Research Center, Hualien Tzu Chi Hospital, Buddhist Tzu Chi Medical Foundation, 707, Section 3, Chung-Yang Road, Hualien, Taiwan.
 E-mail: cyhuang@mail.cmu.edu.tw

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Ho TJ, Ahmed T, Shibu MA, Lin YJ, Shih CY, Lin PY, et al. A prospective review of the health-promoting potential of Jing Si Herbal Tea. Tzu Chi Med J 2024;36(1):1-22.

As CAM use has expanded dramatically, medical professionals should inquire about patients' CAM use concurrently with their medical history. In addition to patients' interest in CAM, researchers have also explored the effectiveness of various therapies and interventions [2].

Traditional Chinese medicine (TCM), an essential category of CAM, is renowned for Chinese herbal medicine and is gaining popularity in many countries. TCM clinics are frequented both by patients looking for therapy for various conditions and those looking to complement, alter, or substitute Western medicine (WM) [3].

Diseases were originally treated using single herbs. However, researchers gained more clinical expertise over time and recognized that ailments were caused by imbalances between the body's various systems. Therefore, to support the dynamic balance between body systems and achieve the best curative results with the most negligible toxicological impact, researchers now prefer the combination of several herbs based on disease severity [4].

Mirroring the pharmaceutical industry, combining TCM to develop new herbal beverages, commonly known as teas, has gained popularity among health-conscious consumers. In addition, herbal tea is a traditional beverage in many cultures around the globe. In general, herbal beverages are composed of natural components derived from various plant parts, including buds, flowers, roots, stems, leaves, and fruits. Scientific evidence shows that herbal teas are rich sources of natural bioactive compounds. These compounds render various biological effects such as anti-viral, anti-oxidant, anti-bacterial, anti-inflammatory, anti-allergic, vasodilatory, anti-mutagenic, anti-carcinogenic, and anti-aging effects, among others [4]. Considering the pharmacological advantages of integrated TCM, the Buddhist Tzu Chi Medical Foundation of Taiwan has recently developed a novel TCM herbal formula known as Jing Si herbal tea (JSHT) by combining eight herbs [5-7]. JSHT is a collective effort to offer TCM as a solution to help citizens improve their overall health and regain balance of the mind and body [5]. The ingredients of JSHT have been reported to exert anti-COVID, anti-aging, and anti-cancer effects, as well as protection against cardiovascular, metabolic, neural, digestive, and genitourinary disease. In the last few decades, substantial studies have been conducted on the effects of JSHT ingredients against different common human diseases. Nevertheless, studies to determine the therapeutic effects of these ingredients are ongoing. However, a review of the fundamental to clinical studies on the potential health benefits of JSHT and its ingredients on various human diseases may provide a useful overview to guide decisions for both consumers and the scientific community. To better understand the beneficial effects of the ingredients in JSHT on health, we searched for relevant articles on *Google Scholar*, *Web of Science*, *PubMed*, *NCBI*, *Science Direct*, *ResearchGate*, and *clinicaltrials.gov*. This review first details the individual ingredients of JSHT and their major phytochemical compounds then summarizes its possible therapeutic effects and discusses the mechanism of action

and toxicity. In light of this, we hope this review can provide a new foundation for further research on JSHT.

INGREDIENTS OF JSHT AND THEIR MAJOR PHYTOCHEMICAL COMPOUNDS

At the end of 2020, Hualien Tzu Chi Hospital developed JSHT for promoting health and wellness. JSHT, approved by the Taiwan Ministry of Health and Welfare, is the brainchild of a joint effort between WM and TCM professionals. JSHT was prepared with selected local Taiwanese herbs, including leaves of *Anisomeles indica* (L.) Kuntze, *Artemisia argyi*, *Ophiopogon japonicus*, and *Perilla frutescens*; roots of *Houttuynia cordata*, *Glycyrrhiza glabra*, and *Platycodon grandifloras*; and flowers of *Chrysanthemum morifolium* Ramat. Different authors have extensively investigated the major chemical constituents of each ingredient of JSHT and their diverse bioactivity. Therefore, this section introduces and describes in detail the various ingredients of JSHT and their major phytochemical compounds [Figure 1 and Table 1].

A. indica (L.) Kuntze (family: *Lamiaceae*) is an essential plant for ethnomedicine, with a long history of application in the treatment of various ailments within the context of TCM. The plant can be found across tropical regions, particularly in Southeast Asia, including Taiwan, China, India, Vietnam, Indonesia, Thailand, Philippines, and Australia. The entirety of the plant has a variety of medical applications and currently, multiple phytochemicals have been prepared from the crude extracts of various parts of the plant. The plant contains flavones, alkaloids, arachnoids, and terpenoids such as ovatodiolide, anisomelic acid, iso-ovatodiolide, 4,7-oxy-cycloanisomelic acid, stigmaterol, and β -sitosterol. The major phytochemicals found in the leaves include tannin, alkaloids, carotenoids, glycosides, and saponin. The seeds contain stigmaterol, β -sitosteroltetracosine, macrocyclic diterpenes, and β -tetracornel- β -amyrin while the flowers contain ovatodiolide, anisomelic acid, and macrocyclic diterpenes [16].

A. argyi (family: *Asteraceae*) is a plant endemic to the northern temperate regions, particularly in Asia. In East Asia, it has a long history of being used as a culinary plant and dietary supplement. *A. argyi* contains a wide range of nutrients, including minerals, polyunsaturated fatty acids, proteins, essential amino acids, dietary fibers, and numerous bioactive compounds such as polysaccharides, flavonoids, organic acids, essential oil, and coumarins. *A. argyi* also contains many essential phytochemical compounds—namely eupatilin, jaceosidin, 5,6-dihydroxy-7,3,4-trimethoxyflavone, 5,6,4-trihydroxy-7,3-dimethoxyflavone, 7-O- β -D-glucopyranoside, 5,6,2,4-tetrahydroxy-7,5-dimethoxyflavone, artemisian A-D, isoartemisolid/sesquiterpene dimer (DSF-52), 3,5-dicaffeoylquinic acid, scopoletin, terpenoids, alcohols, ethers, and ketones [17].

C. morifolium Ramat has a wide range of applications, including effective treatments for various ailments in traditional medicine and nutritious herbal tea in the food industry. These perennial plants are prevalent in Asia and northeastern Europe, with most species being indigenous



Figure 1: The main ingredients of Jing Si herbal tea. Image of ingredients adopted from [8-15].

to East Asia. *Chrysanthemum* plants belong to the *Asteraceae* family, which contains low-molecular-weight components, including flavonoids, triterpenes, unsaturated fatty acids, and sesquiterpenes. *Chrysanthemum* spp. leaves contain chlorogenic acid, daucosterol, β -sitosterol, octa-cosyl alcohol, lupeol, ineupatorolide B, syringin, petasiphenol, physcion, acacetin, eupatilin, quercetin, diosmetin, luteolin, apigenin, apigenin-7-O- β -D-glucopyranoside, quercetin-3-O- β -D-glucopyranoside, luteolin-7-O- β -D-glucopyranoside, apigenin-7-O- β -D-neosphaeroside, and acacetin-7-O- β -D-glucoside. In addition, most *Chrysanthemum* spp. flowers contain anthocyanins, cyanidin 3-glucoside and cyanidin 3-(3''-malonyl) glucoside, and carotenoids—lutein, zeaxanthin, β -cryptoxanthin, 13-cis- β -carotene, trans- β -carotene, and 9-cis- β -carotene [18].

G. glabra belongs to the family Fabaceae (also known as Leguminosae) and has a variety of medicinal uses. Members of this family are also commonly used as feed and food. This species is native to Mediterranean areas but is now also present in Taiwan, China, Russia, and India. Recently, a large number of biological compounds have also been isolated from *G. glabra*, mostly triterpenes, saponins, and flavonoids. Glycyrrhizin, a triterpenoid saponin approximately 50 times sweeter than sucrose, is the principal active component of the roots. The major flavonoids are glycosides of isoliquiritigenin (2',4,4'- trihydroxychalcone) and liquiritigenin (4',7-dihydroxyflavanone) such as liquiritin

apioside, liquiritin, licuraside, and isoliquiritin. Moreover, five new flavonoids have been isolated from the dried roots: Apioside, glucoliquiritin, shinpterocarpin, shinflavanone, 1-methoxyphaseolin, and prenyllicoflavone A. In addition, licoflavanone and pinocembrin were also isolated from the leaves [19].

H. cordata, a perennial herb belonging to the Saururaceae family, is widely utilized in both Chinese herbal medicine and for food. It prefers moist soil and warm conditions for growth. Its applicability in Taiwan, China, Japan, Korea, India, and other Asian countries has been described. Over the past few 1000 years, local people have eaten as well as used *H. cordata* as medicine, and it is still used as food and medicine. *H. cordata* has several chemical components, with alkaloids being the most abundant; among them, most are phenanthrolactam compounds such as aristolactam and piperolactam. Moreover, *H. cordata* flavonoids include hyperoside, rutin, quercitrin, and quercetin, most of which are glycosides containing rhamnose. However, phenolic acids – including linolenic, stearic, palmitic, oleic, chlorogenic, cryptochlorogenic, and neochlorogenic acid and caffeic and quinic acid derivatives, and other ingredients – are the most isolated components [20].

Ophiopogon japonicus (*Ophiopogon* genus, Liliaceae family) is both a food and traditional herbal medicine in Taiwan, China, Japan, and some south-eastern Asian

Table 1: Major phytochemical compounds of the ingredients of Jing Si herbal tea

Ingredients	Major phytochemical compounds	Reference
<i>A. indica</i> (L.) Kuntze	Ovatodioid, Alpha-Humulene, 2TCH2H, 4DHNT, Stigmasterol, Bicyclo [8.1.0] undecane, 2-(4-Methylphenyl) ethynylaniline, Vitamin E, E12TD	[16]
<i>A. argyi</i>	Eupitalin, Jaceosidin, 5,6-dihydroxy-7,3,4-trimethoxyflavone, 5,6,4-trihydroxy-7,3-dimethoxyflavone, 7-O-β-D-glucopyranoside, 5,6,2,4-tetrahydroxy-7,5-dimethoxyflavone, Artemisian A-D, Isoartemisolid/DSF-52, 3,5-dicaffeoylquinic acid, Scopoletin, Terpenoids, Alcohols, Ethers, Ketones	[17]
<i>C. morifolium</i> Ramat	Chlorogenic Acid, Luteolin-7-O-β-Glucoside, 3,5-Dicaffeoyl Quinic Acid, Apigenin-7-O-β-Glucoside, Linarin, Acacetin-7-O-β-Glucoside, Luteolin, Apigenin, Cyanidin-3-O-Glucoside, Cyanidin-3-O-(6-Malonyl)Glucoside, Cyanidin, Carotenoids, dendranlignan A	[18]
<i>G. glabra</i>	Liquiritin, Isoliquiritin, Glycyrrhizin, Liquiritigenin, Isoliquiritigenin, 18-β-Glycyrrhetic acid, Liquiritin apioside, Glycyrrhetic acid, Licochalcone A, Glabridin, Glycycomarin	[19]
<i>H. cordata</i>	Sodium houttuynfonate, Houttuynin, Houttuynoid, Houttuynoside, Bornyl acetate, β-myrcene, Ethyl caprate, Ethyl dodecanoate, 2-undecanone, 2-tridecanone, Quercetin, Linolenic acid, Oleic acid, Palmitic acid, Stearic acid, Neochlorogenic acid, Chlorogenic acid, 4-dicaffeoylquinic acid, Aristololactam, Rutinum, Hyperoside	[20]
<i>O. japonicus</i>	Saponins, DT-13, Ophiopogonin, Ophiopogonide, Ophiopogonane, Ruscogenin, β-d-fructan, FOJ-5, Methylphiopogona, Nolinospinoside, Pulullan, lectin, liposomes, 8-Formylophiopogonanone B	[21]
<i>P. frutescens</i>	Phenolic acids (rosmarinic acid, rosmarinic acid-3-O-glucoside, caffeic acid, caffeic acid-3-O-glucoside, ferulic acid), Flavonoids (catechin, apigenin, apigenin 7-O-glucuronide, apigenin 7-O-diglucuronide, luteolin, luteolin 7-O-glucuronide, luteolin 7-O-diglucuronide, scutellarein, scutellarein 7-O-glucuronide, scutellarein 7-O-diglucuronide), Anthocyanins (shisonin, malonylshisonin, cyanidin 3-O-caffeoylglucoside-5-O-glucoside, cyanidin 3-O-caffeoylglucoside-5-O-malonylglucoside), triterpene acids, Fatty acids (palmitic acid, stearic acid, lauric acid, oleic acid, linoleic acid, linolenic acid), Tocopherols (α-tocopherol, β-tocopherol, γ-tocopherol, δ-tocopherol), Phytosterols (campesterol, stigmasterol, β-sitosterol, β-amyirin), Perillaldehyde	[22]
<i>P. grandiflorus</i>	Triterpenoid saponins, platyconin, taxifolin, platycoside, quercetin-7-O-glucoside, quercetin-7-O-rutinoside, luteolin-7-O-glucoside, apigenin-7-O-glucoside, luteolin, apigenin, palmitic, oleic acids, caffeic acid, 3,4-dimethoxycinnamic acid, ferulic acid, isoferulic acid, m-coumaric acid, p-coumaric acid, p-hydroxybenzoic acid, aresorcylic acid, 2,3-dihydroxybenzoic acid, 2-hydroxy-4-methoxybenzoic acid, homovanillic acid, chlorogenic acid, lobetylol, lobetyolin, lobetyolinin, spinasterol, α-spinasteryl-3-O-β-D-glucoside, betulin, β-sitosterol, and δ-7-stigmastene-3, platycodin D	[23]

2TCH2H: 2,6,10,14,18,22-Tetracosahexaene, 2,6,10,15,19,23-hexamethyl-, (all-E)-, 4DHNT: 4-Dehydroxy-N-(4,5-methylenedioxy-2-nitrobenzylidene) tyramine, E12TD: Ethane, 1-(4,4,4-trifluoro-1,3-dithiobutyl)-2-(3,3,3-trifluoro-1,2-dithiopropyl), DT-13: Ruscogenin 1-O-[[β-D-glucopyranosyl-(1→2)]-β-D-xylopyranosyl-(1→3)]-β-D-fucopyranoside, FOJ-5: Fructosan with an average molecular weight of 5 kDa from *O. japonicus*, *A. indica*: *Anisomeles indica*, *A. argyi*: *Artemisia argyi*, *C. morifolium*: *Chrysanthemum morifolium*, *G. glabra*: *Glycyrrhiza glabra*, *H. cordata*: *Houttuynia cordata*, *O. japonicus*: *Ophiopogon japonicus*, *P. frutescens*: *Perilla frutescens*, *P. grandiflorus*: *Platycodon grandiflorus*

countries. Roots of *O. japonicas* are considered an effective clinical tonic and strengthening drug for treating various disorders. Several substances, including saponins, DT-13 (ruscogenin 1-O-[[β-D-glucopyranosyl-(1→2)]-β-D-xylopyranosyl-(1→3)]-β-D-fucopyranoside), ophiopogonin, ophiopogonide, ophiopogonane, ruscogenin, β-d-fructan, FOJ-5 (fructosan with an average molecular weight of 5 kDa), methylphiopogonanone A/B, nolinospinoside, pulullan, lectin, liposomes, and 8-formylophiopogonanone B, have been identified from various parts of *Ophiopogon japonicus*. The primary active components of *Ophiopogon japonicus* are steroidal saponins and homoisoflavonoids, which have various pharmacological actions [21].

P. frutescens is an annual herbaceous, aromatic, edible, and attractive plant, a member of the Lamiaceae family. In East Asian countries (Taiwan, China, Korea, Japan, India, and Vietnam), it has been used as a culinary and medicinal herb for thousands of years. In addition, folk medicine uses the leaves, seeds, and stems of *P. frutescens* for a variety of therapeutic purposes. Perilla seeds, stems, and leaves contain 271 identified and documented phytochemical substances. These active compounds may be categorized either as hydrophilic (flavonoids, anthocyanins, and phenolic acids) or lipophilic (volatile compounds,

policosanols, tocopherols, triterpenes, phytosterols, and fatty acids). The primarily identified phytochemicals are phenolic acids (rosmarinic acid: Rosmarinic acid-3-O-glucoside; caffeic acid: Caffeic acid-3-O-glucoside; and ferulic acid), flavonoids (catechin, apigenin, apigenin 7-O-glucuronide, apigenin 7-O-diglucuronide, luteolin, luteolin 7-O-glucuronide, luteolin 7-O-diglucuronide, scutellarein, scutellarein 7-O-glucuronide, and scutellarein 7-O-diglucuronide), anthocyanins (shisonin, malonylshisonin, cyanidin 3-O-caffeoylglucoside-5-O-glucoside, and cyanidin 3-O-caffeoylglucoside-5-O-malonylglucoside), triterpene acids, fatty acids (palmitic, stearic, lauric, oleic, linoleic, and linolenic acid), tocopherols (α-, β-, γ-, and δ-tocopherol), phytosterols (campesterol, stigmasterol, β-sitosterol, and β-amyirin), and perillaldehyde [22].

Platycodon grandiflorus, the sole species in the family Campanulaceae, has long been used conventionally as herbal medicine. In addition to its usage as a legal medicine and nutritional supplement, it is a common element in healthy foods and vegetable meals in Northeast Asia (including Taiwan, China, Korea, and Japan). Natural products extracted from *Platycodon grandiflorus* can be classified into different classes and display different structural characteristics. In the past few decades, at least 100

compounds have been isolated from *Platycodon grandiflorus*, including triterpenoid saponins, platyconin, taxifolin, platycoside, quercetin-7-O-glucoside, quercetin-7-O-rutinoside, luteolin-7-O-glucoside, apigenin-7-O-glucoside, luteolin, apigenin, palmitic, oleic acids, caffeic acid, 3,4-dimethoxycinnamic acid, ferulic acid, isoferulic acid, m-coumaric acid, p-coumaric acid, p-hydroxybenzoic acid, aresorcylic acid, 2,3-dihydroxybenzoic acid, 2-hydroxy-4-methoxybenzoic acid, homovanillic acid, chlorogenic acid, lobetyol, lobetyolin, lobetyolinin, spinasterol, α -spinasteryl-3-O- β -D-glucoside, betulin, β -sitosterol, and δ -7-stigmastene-3, and platycodin D [23].

THERAPEUTIC EFFECTS OF JSHT AND ITS INGREDIENTS

The ingredients of JSHT have been documented to have multiple biological impacts on health promotion and treatment of various diseases. Several studies have reported various pharmacological activities, including anti-COVID, -aging, -cancer, -diabetic, -obesity, and -inflammatory activities, and cardio-, hepato-, nephro-, and neuroprotective effects [Figure 2]. To explain the nature of individual JSHT ingredients, this review intends to summarize their pharmacological and therapeutic effects and potential mechanisms against various disorders based on the relevant research for future clinical use and experimental reference.

Anti-COVID-19 activity

Since the outbreak of COVID-19, health-care professionals worldwide have concentrated on devising methods to combat the virus, and TCM experts have joined the endeavor. Therefore, several anti-coronavirus health products based on TCM formulas have arrived in the market due to the latter's efforts. At the end of 2020, Hualien Tzu Chi Hospital developed JSHT for health and wellness. In a clinical trial, researchers from the Buddhist Tzu Chi Medical Foundation investigated the efficacy and safety of JSHT combined with standard treatment for patients with mild-to-moderate COVID-19 [Table 2] [5]. The results demonstrated that in adult patients with mild-to-moderate COVID-19, the combination of JSHT and standard therapy significantly improved C-reactive protein level, reverse transcription-polymerase chain reaction

threshold value, and Brixia score—most notably in males and older patients (aged >60). The patients also experienced 51, 70, and 100% lower risks of intubation, Medisave Care Unit admissions, and mortality, respectively, compared to those receiving standard management alone. This suggests that supplemental JSHT treatment can avert critical status and fatality. Li *et al.* [30] performed a retrospective cohort study on severe COVID-19 patients to investigate the impact of JSHT-incorporated treatment strategy on death and hospital course. The study recruited 10 patients divided into two groups: >70 years and <70 years and each group contained five patients. In all patients, JSHT was co-treated with death as the primary outcome. Laboratory tests, computed tomography (CT) values, clinical course, and hospital stays were considered secondary outcomes. Study results indicate that older patients (>70 years) have higher Charlson Comorbidity Index scores and Veterans Health Administration COVID-19 scores, as well as lower hemoglobin levels than <70 years old. Moreover, CT values, lymphocyte count, lactate dehydrogenase, and D-dimer were inconsistent among nonsurvivors. The death rate was 20%, and the recovery rate for mild illness in 14 days was 40%. Another research team from the Buddhist Tzu Chi Medical Foundation reported that JSHT demonstrated anti-viral activity in Calu-3 and Caco-2 cells and mice pretreated for 3 days when exposed to distinct severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) variants [6]. Intriguingly, JSHT significantly inhibited 3cL and RdRp activity, demonstrating the multitargeting nature of JSHT and making it a possible SARS-CoV-2 infection preventive. In addition, numerous investigations have demonstrated the anti-viral activity of JSHT's ingredients during the past decade. Recently, a study showed the potent anti-inflammatory properties of *A. indica* (L.) Kuntze may alleviate the uncontrolled release of pro-inflammatory cytokines (NO, tumor necrosis factor [TNF]- α , and interleukin (IL)-12) in response to COVID-19 infection [31]. Similarly, *A. argyi* may have therapeutic potential for decreasing COVID-19-induced inflammatory-related conditions because of its anti-inflammatory capabilities, which are mediated by inhibiting the nuclear factor kappa B (NF- κ B) pathway [32]. *C. morifolium* Ramat contains the




Figure 2: Various pharmacological activities such as anti-covid [24], anti-aging [25], anti-cancer [26], neuroprotective [27], liver and kidney [28], immunomodulator and other effects [29] of the individual ingredients in Jing Si herbal tea

Table 2: Anti-COVID and anti-aging effects of the Jing Si herbal tea and its ingredients

Property	Sample/ ingredients	Experimental model	Outcomes	Mode of action	Reference
Anti-COVID effects	Jing Si herbal tea	<i>In vitro</i> and animal model	Inhibited viral replication	Inhibited viral entry into Calu-3 and Caco-2 cells. Decreased 3cL and RdRp activity	[6]
	-	Clinical study	Reduced the SARS-CoV-2 viral load and systemic inflammation and alleviated lung infiltrates in the patients with mild-to-moderate COVID-19	Improved the reverse transcription-polymerase chain reaction cycle threshold value, C-reactive protein level, and Brixia score	[5]
	-	Clinical study	The death rate was 20% and recovery rate to mild illness in 14 days was 40%	Older patients (>70 years) have higher Charlson Comorbidity Index scores and VACO scores as well as lower hemoglobin levels than <70 years patients	[30]
	<i>A. indica</i> (L.) Kuntze	Animal model	Inhibited pro-inflammatory mediators and tumor cell proliferation	Inhibited NO, TNF- α and IL-12	[31]
	<i>A. argyi</i>	Animal model	Attenuated airway inflammation	Decreased iNOS expression and NF- κ B phosphorylation	[32]
	<i>C. morifolium</i> Ramat	<i>In vitro</i> and animal model	Showed anti-platelet and anti-thrombosis activity	Suppressed TXA2, NF- κ B, and iNOS. Decreased serum levels of TNF- α , IL-1 β , IL-6, and CRP and reduced macrophages, neutrophils, and white blood cells	[33]
	<i>G. glabra</i>	<i>In vitro</i>	Protected against coronavirus	Replication of the virus was inhibited by inducing oxide synthase in Vero cells	[34]
	<i>H. cordata</i>	Bedside-to-bench (<i>in vitro</i> and clinical trial)	Inhibited SARS-CoV-2 pathogenesis in anti-viral and -inflammatory assays	Block binding between ACE2 and spike protein of SARS-CoV-2	[35]
	<i>O. japonicus</i>	Animal model	Inhibited venous thrombosis	Secured endothelial cells from anoxic injury and decreased leukocytes adhesion and inflammation of the vein wall	[36]
	<i>P. frutescens</i>	<i>In vitro</i>	Inhibited SARS-CoV-2 replication by inactivating the virion	Inactivate SARS-CoV-2 spike protein Decrease viral protein/RNA synthesis and the levels of TNF- α , IL-1, and IL-6	[37]
Anti-aging effects	<i>P. grandiflorus</i>	<i>In vitro</i>	Platycodin D showed a protective effect against SARS-CoV-2 infection	Blocked lysosome and TMPRSS2-driven viral entry	[38]
	Jing Si herbal drink	<i>In vitro</i> and animal model	Displayed pleiotropic effects against aging-associated disorders	Maintained stem cell homeostasis and provided cytoprotection. Regulated blood glucose metabolism, enhanced neurons' autophagic clearances, and suppressed cancer growth and migration	[25]
	<i>A. indica</i> (L.) Kuntze	<i>In vitro</i>	Ovatodiolide compound showed inhibitory effects on melanogenesis	Inhibited mushroom tyrosinase activity (IC ₅₀ =0.253 mM), suppressed intracellular tyrosinase activity (IC ₅₀ =0.469 mM), and decreased the amount of melanin (IC ₅₀ =0.435 mM)	[39]
	<i>A. argyi</i>	Animal model	Exerted ameliorative effects on DNCB-induced ADlike lesions by exerting antiallergenic skin inflammatory effects related to the recovery of skin barrier dysfunction	Suppressed cytokine abundance via the regulation of crucial factors, including Lyn, Syk, MAPKs, PI3K/Akt, and I κ B α , during the process of AD pathogenesis	[40]
	<i>C. morifolium</i> Ramat	Clinical study	Significantly improved erythema and overall rosacea severity in patients with moderate rosacea, with mild adverse reactions observed in both the treatment and placebo groups		[41]
	-	Clinical study	Inhibited mushroom tyrosinase activity. Reduced facial melanin levels. No moisture, elasticity, wrinkles, evenness, and pore size changes		[42]
	<i>G. glabra</i>	Clinical study	Lightened hand solar lentigines		[43]

Contd...

Table 2: Contd...

Property	Sample/ ingredients	Experimental model	Outcomes	Mode of action	Reference
	<i>H. cordata</i>	<i>In vitro</i>	Inhibited UVB-irradiated skin aging through regulation of the MAPK signaling pathway by attenuating the activation of JNK/ERK/c-Jun in human dermal fibroblasts	Inhibited intracellular ROS production and inflammatory cytokine secretions (IL-6 and IL-8) while increasing collagen type I synthesis along with downregulating MMP-1 gene and protein expressions	[44]
	<i>O. japonicus</i>	Animal model	Extended lifespan in <i>C. elegans</i>	Slightly increased the pharyngeal pumping rate of <i>C. elegans</i> and reduced the accumulation of lipofuscin	[45]
	<i>P. frutescens</i>	<i>In vitro</i> and clinical study	Improved skin elasticity and reduced skin pigmentation	Reduced retraction time and melanin content	[46]
	-	Clinical study	Increased skin hydration and skin elasticity	The sulcus cutis was narrower with the effect of the test serum, and the number of equilateral triangles was increased than the baseline	[47]
	<i>P. grandiflorus</i>	Animal model	Alleviated DNCB-induced atopy-like dermatitis	Reduced the elevation of serum total IgE or cytokine mRNA. Suppressed Th1 and Th2 responses	[48]

VACO: Veterans Health Administration COVID-19, ACE2: Angiotensin-converting enzyme 2, IL-1 β : Interleukin 1 beta, IL-6: Interleukin 6, iNOS: Inducible nitric oxide synthase, TXA2: Thromboxane A2, NF- κ B: Nuclear factor kappa B, NO: Nitric oxide, RdRP: RNA dependent RNA polymerase, TNF: Tumor necrosis factor, DNCB: 1-chloro-2, 4-dinitro benzene, AD: Atopic dermatitis, Lyn: Lck/Yes novel tyrosine kinase, SyK: Spleen tyrosine kinase, MAPKs: Mitogen-activated protein kinases, PI3K/Akt: Phosphatidylinositol-3-kinase and protein kinase B, I κ B α : Nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, alpha, UVB: Ultraviolet B, JNK: Jun N-terminal kinase, ERK: extracellular regulated kinase, ROS: Reactive oxygen species, MMP-1: Matrix metalloproteinase-1, IgE: Immunoglobulin E, mRNA: Messenger RNA, Th1: T helper cell type 1, Th2: T helper cell type 2, SARS-CoV-2: Severe acute respiratory syndrome coronavirus 2, CRP: C-reactive protein, IC₅₀: Half maximal inhibitory concentration, *C. elegans*: *Caenorhabditis elegans*, *A. indica*: *Anisomeles indica*, *A. argyi*: *Artemisia argyi*, *C. morifolium*: *Chrysanthemum morifolium*, *G. glabra*: *Glycyrrhiza glabra*, *H. cordata*: *Houttuynia cordata*, *O. japonicus*: *Ophiopogon japonicus*, *P. frutescens*: *Perilla frutescens*, *P. grandiflorus*: *Platycodon grandiflorus*

flavonoids luteolin and apigenin, which have antithrombotic properties. These compounds effectively suppressed the production of thromboxane A2 and collagen-induced platelet aggregation in an *in vitro* human investigation, suggesting their potential use in treating or preventing blood clots [33]. Moreover, casticin from *C. morifolium* Ramat, another polymethylflavone, may inhibit tissue damage and cytokine storms during COVID-19 infection by inhibiting the NF- κ B and inducible nitric oxide synthase (iNOS) pathways and reducing serum levels of pro-inflammatory cytokines (TNF- α , IL-1, and IL-6) and C-reactive protein. Further, glycyrrhizic acid, the active component of *G. glabra*, efficiently inhibits the entry and replication of the SARS virus [34]. Furthermore, it can inhibit entry, replication, and inflammation of SARS-CoV-2 by regulating steroid metabolism, blocking the main protein of SARS-CoV-2, and inhibiting the release of inflammatory agents that lead to cytokine storms. Tsai *et al.* [35] devised NRICM101, a new TCM formulation, as a COVID-19 treatment for a bench-to-bedside study. *H. cordata*, one of the constituents of NRICM101, can reportedly block the SARS-CoV-2 spike RBD protein from binding to ACE2 and reduce TNF- α production. COVID-19 mortality is associated with thrombotic complications such as pulmonary emboli, venous thromboembolic disease, and stroke. In a recent study, *O. japonicus* remarkably reduced venous thrombosis [36]. Anoxic injury to endothelial cells was prevented by ethanol extracts of *O. japonicus*, which also reduced leukocyte adhesion to the endothelium and suppressed inflammation of the vein walls in rat and mouse models, further supporting its therapeutic potential for COVID-19. Another recent *in vitro* study using Calu-3 cells showed that *P. frutescens* leaf extract (PLE) could inhibit virus protein/RNA synthesis as well as virus-induced cytokines such as

IL-1, IL-6, and TNF- α [37]. In addition, PLE prevents viral entry by inactivating the SARS-CoV-2 virus, i.e., the spike protein. Notably, the combination of PLE and remdesivir has a synergistic impact against SARS-CoV-2. Platycodin D (a bioactive compound of *P. grandifloras*) is capable of redistributing membrane cholesterol to block the main infectious routes for SARS-CoV-2, including lysosomes and TMPRSS2 [38]. Considering the pharmacological effects mentioned above, JSHT has excellent therapeutic potential to prevent and improve the clinical outcomes of COVID-19. However, due to the lack of studies on human subjects, more clinical trials focused on patient groups, efficacy, and doses are required. Furthermore, the effects of JSHT on anhedonia and depressive symptoms linked to long COVID were assessed in a recent study using mice models subjected to unpredictable chronic mild stress [49]. The study's results demonstrate that JSHT mitigates the cytokine storm by inhibiting NF- κ B cascades, offering protective effects against symptoms connected with long-term COVID-19 infection.

Anti-aging activities

Aging can be defined as the progressive loss of cells, tissues, and organs throughout a person's lifetime. The loss of tissue homeostasis and normal function increases susceptibility to age-related illnesses by reducing adaptation to varied stress circumstances. Consequently, aging is one of the critical risk factors for various disorders. To promote healthy aging, pharmacological interventions must target a variety of factors associated with aging disorders and simultaneously promote adult stem cell rejuvenation. TCM has demonstrated rejuvenation potential in a variety of experimental models. In this context, JSHT and its constituent compounds obtained from natural sources with potential properties, which could achieve similar benefits as synthesized compounds, are of

great interest. In search of a promising pleiotropic medicine for aging treatment, a research team from the Buddhist Tzu Chi medical foundation tested JSHT for its anti-aging benefits using relevant *in vitro* and *in vivo* models [25] [Table 2]. Their findings indicated that JSHT preserved stem cell homeostasis and offered cytoprotection. In addition, it inhibited the growth and migration of cancer cells, promoted autophagic clearing in neurons, and affected blood glucose metabolism. A research team concluded that JSHT is a possible pleiotropic agent for healthy aging since it acts on multiple sites and provides a cumulative protective effect against numerous age-associated diseases. Several other studies also showed the potential anti-aging effects of the ingredients in JSHT. For instance, Huang *et al.* studied the antioxidative properties of *A. indica* (L.) Kuntze methanol extract and the inhibitory effect of its active component, ovatodiolide, on melanogenesis in B16F10 melanoma cells [39]. The results revealed that mushroom tyrosinase activity ($IC_{50} = 0.253$ mM) was inhibited by purified ovatodiolide. In addition, the compound effectively decreased the amount of melanin ($IC_{50} = 0.435$ mM) due to suppressed intracellular tyrosinase activity ($IC_{50} = 0.469$ mM) in a dose-dependent manner. Atopic dermatitis (AD) is a prevalent recurrent inflammatory skin condition. In one study, *A. argyi* extract ameliorated DNCB-induced AD-like lesions by exerting anti-allergenic restorative skin inflammatory actions after skin barrier failure [40]. These effects are thought to be related to suppressing cytokine abundance through regulating various factors involved in AD pathogenesis—including IkB α , Syk, Lyn, MAPKs, and PI3K/Akt. Similarly, *Platycodon grandiflorus* showed potential effects on AD in DNCB-treated NC/Nga mice [48]. Oral treatment of PG is considered to reduce the elevated levels of serum total IgE or cytokine mRNA. *Platycodon grandiflorus* probably alleviates AD-like skin lesions by inhibiting both Th1 and Th2 cytokines. Similar to AD, rosacea is an inflammatory dermatosis with a complex etiology involving systemic or local, endogenous, and environmental factors. Rosacea is considered a disease characterized by vascular changes. As an herbal extract containing a potent combination of saponins, phenylpropenoic acids, and flavonoids, *C. morifolium* Ramat is well-known for improving capillary mechanical resistance and decreasing vascular wall permeability. Clinical trial findings showed that, compared to baseline and placebo, treatment with *C. morifolium* Ramat extract-based cream significantly reduced the rosacea and erythema severity and overall severity efficacy assessment scores ($P = 0.046$ and $P = 0.001$, respectively) [41]. In addition, another clinical study assessed the effects of *C. morifolium* Ramat extract in cosmetic formulations on biophysical skin parameters in healthy people [42]. Dose-dependent inhibition of tyrosinase activity was observed with *C. morifolium* Ramat extracts. Furthermore, time-dependent reductions in facial melanin levels were also observed in the faces of human volunteers who were exposed to daily cosmetic formulations with 0.5% *C. morifolium* Ramat water extract. Benign hyperpigmentation, another age-related, often occurs on sun-exposed skin in adults (face, hands) due to photoaging, postinflammatory, or chronoaging processes. Age-related blemishes are caused by a limited, benign proliferation of melanocytes at the

dermo-epidermal interface, producing aesthetic issues. Grippaudo and Di Russo [43] conducted a clinical trial to evaluate the efficacy of glycyrrhetic acid (extracted from *G. glabra*) monotherapy combined with fractional CO₂ laser for benign hand hyperpigmentation treatment. The hand solar lentigines of all participants, investigators, and blinded dermatologists improved. *P. frutescens*, another ingredient of JSHT, can effectively reduce the effects of skin aging and hyperpigmentation disorders. An *in vitro* and clinical study by Mungmai *et al.* [46] revealed that the anti-melanogenic effects of *P. frutescens* on B16F10 cells occur without cytotoxicity or death. In addition although skin elasticity improved slightly in the 4th week compared to the third, it was found that the amount of melanin in the skin significantly decreased ($P < 0.05$) without any irritation in the 4th week. The same research team conducted another clinical trial on the efficacy of cosmetic formulation containing PLE (PLES) for aging skin [47]. No skin irritation or allergic reaction was reported during application and following PLES treatment, skin hydration and elasticity could increase. However, frequent exposure to UV radiation generates changes in an individual's skin characterized by tanning, burning, elastin fibers, loosening collagen, and diminished skin integrity, known as photoaging. A fraction enriched in hyperosides produced by *H. cordata* Thunb was tested for photoprotection against UVB-induced aging of human fibroblasts [44]. Activation of JNK/ERK/c-Jun in human dermal fibroblasts was attenuated by the hyperoside-enriched fraction obtained from *H. cordata*. Recently, there has been an increasing interest in *O. japonicas* as an anti-aging agent. Yu *et al.* [45] reported that *O. japonicas* is capable of extending the lifetime of nematodes as well as reversing the age-related drop in pharyngeal pumping and decreasing the formation of the age pigment lipofuscin in the animals.

Anti-cancer activities

Cancer is a complex disease characterized by either increased or decreased cell proliferation, resulting in apoptosis. Exogenous and endogenous causes might contribute to excessive reactive oxygen species (ROS) generation. Thus, DNA is broken or mutated, chromosomes rearranged, DNA cross-links formed, nucleic acids degrade, the cell membrane becomes damaged due to lipid peroxidation, and tumors can form. JSHT ingredients have shown anti-cancer properties in both *in vitro* and *in vivo* studies [Table 3]. Ovatodiolide, which is present in the plant *A. indica* (L.) Kuntze, reportedly has an anti-proliferative effect on several cancer cells. In AGS cells, the pure ovatodiolide reduced cell proliferation and triggered cell death in a dose-dependent manner, accompanied by G2/M cell cycle arrest and nuclear condensation and fragmentation [50]. Activation of Bax and reduction in Bcl-2 mRNA levels could be potential causes, at the very least in part, of the apoptotic effects of ovatodiolide on AGS cells. Moreover, ovatodiolide can effectively suppress oral tumorigenesis and stemness properties through JAK2/STAT3 signaling [51]. *A. argyi* polysaccharide fraction reportedly prevents tumor cell-induced platelet aggregation by inhibiting the interaction of podoplanin with C-type lectin-like receptor 2 [52]. Furthermore, computational tools

Table 3: Anti-cancer and cardiovascular protective effects of the ingredients of Jing Si herbal tea

Property	Sample/ ingredients	Experimental model	Outcomes	Mode of action	Reference
Anti-cancer	<i>A. indica</i> (L.) Kuntze	<i>In vitro</i>	Ovatodiolide compound significantly inhibited cell proliferation and induced cell apoptosis of human gastric cancer cells	Activated <i>Bax</i> and the reduction in <i>Bcl-2</i> mRNA levels	[50]
	-	<i>In vitro</i>	Ovatodiolide effectively suppressed oral tumorigenesis and stemness properties of oral cancer stem-like cells	Suppressed oral-sphere formation, reduced clonogenicity, enhanced cisplatin sensitivity and dysregulated JAK2/STAT3 signaling pathway	[51]
	<i>A. argyi</i>	<i>In vitro</i>	Prevention of tumor cell metastases	Suppressed PDPN- and tumor cell-induced platelet aggregation by irreversibly blocking the interaction between PDPN and CLEC-2	[52]
	<i>C. morifolium</i> Ramat	<i>In vitro</i>	Inhibited the proliferation of MCF7 cells	Regulates the biological processes of their protein targets	[53]
	<i>G. glabra</i>	<i>In vitro</i>	18 β -glycyrrhetic acid potently inhibited the migration and invasion of colorectal cancer cells and reduced MMP expression as well	Suppressed PI3K and STAT3 signaling pathways	[54]
	-	<i>In vitro</i> and animal model	Glycycomarin exerted anti-liver cancer activity	Inactivated oncogenic kinase TOPK, which in turn led to activation of its binding partner p53, followed by cell cycle arrest and cell death induction	[55]
	-	Animal model	Glycyrrhizin exerted anti-cancer action in lung cancer cell lines	Suppressed the TxA2 pathway	[56]
	<i>H. cordata</i>	<i>In vitro</i>	Showed anti-carcinogenic activity against breast cancer	Induced cell cycle arrest via the modulation of cyclin D1, CDK4, and p21 protein expressions. Suppressed cell migration and invasion by inhibiting MMP-1 and MMP-9 secretion	[57]
	-	<i>In vitro</i>	Induced apoptosis in human HepG2 hepatocellular carcinoma cells	Promoted the activation of HIF-1 α -FOXO3 and MEF2A pathways	[58]
	<i>O. japonicus</i>	<i>In vitro</i>	Ophiopogonin-B decreased proliferation and induced apoptosis of human gastric cancer cells	Anti-cancer mechanisms are associated with the JNK1/2 and ERK1/2 signaling pathways	[59]
	-		O-D inhibited cell proliferation and induced apoptosis in human breast carcinoma MCF-7 cells	O-D-induced G ₂ /M cell cycle arrest was associated with down-regulation of cyclin B1. Activation of caspase-8 and caspase-9 was associated with ophiopogonin D-induced apoptosis	[60]
	<i>P. frutescens</i>	<i>In vitro</i> and animal model	Synergistically suppressed proliferation of A549 human lung adenocarcinoma	Induced two-stage cell cycle arrest at G ₁ and G ₂ /M phases	[61]
	<i>P. grandiflorus</i>	<i>In vitro</i>	Induced cell apoptosis and inhibited cell proliferation	Inhibited the expression of p-Akt and p-STAT3	[62]
	-	<i>In vitro</i>	Induced cancer cell death is associated with cytoplasmic pinocytic and autophagic vacuolation	Activated AMP-activated protein kinase	[63]
Cardioprotective	<i>A. argyi</i>	Animal model	Showed antihypertensive activity in spontaneously hypertensive rats	The underlying mechanism appears to involve, at least in part, an increase in urine and electrolyte output	[64]
	<i>C. morifolium</i> Ramat	Animal model	Alleviated hypertensive cardiac hypertrophy in rats	Inhibited HIF-1 α expression and modulated PPAR α -mediated CPT-1 α , PDK-4, and GLUT-4 expressions	[65]
	<i>G. glabra</i>	Animal model	18 β -glycyrrhetic acid showed cardioprotective effects on acute myocardial infarction	Inhibited oxidative stress, inflammation, apoptosis via the PI3K/Akt pathway, and reduced cell contractility and Ca ²⁺ concentration via L-type Ca ²⁺ channels	[7]

Contd...

Table 3: Contd...

Property	Sample/ingredients	Experimental model	Outcomes	Mode of action	Reference
-		<i>In vitro</i> and animal model	<i>G. glabra</i> extracts showed cardioprotective effects against doxorubicin-induced cardiotoxicity	Decreased serum LDH and CK-MB levels, improved heart morphology, and increased GSH-PX activity and GSH level	[66]
	<i>H. cordata</i>	<i>In vitro</i> and animal model	2-undecanone protected against fine particle-induced heart inflammation	Modulated Nrf2/HO-1 and NF-κB pathways	[67]
	<i>O. japonicus</i>	Animal model	8-Formylophiopogonanone B antagonized doxorubicin-induced cardiotoxicity	Suppressed heme oxygenase-1-dependent myocardial inflammation and fibrosis	[68]
	<i>P. frutescens</i>	<i>In vitro</i>	Protected endothelium against vascular inflammation	Inhibited ICAM-1 expression and IL-6 production, adherence of leukocytes to endothelial cells, and EMPs generation	[69]
-		Animal model	Perillaldehyde prevented the formation of atherosclerotic plaques	Increased tetrahydrobiopterin levels and subsequent eNOS recouping	[70]
	<i>P. grandiflorus</i>	Animal model	Platycodin D provided myocardial protection in spontaneously hypertensive rats	Downregulated the expression of FGF2, uPA, MMP2, MMP9, and CTGF and decreased TGF-β1 expression	[71]

Bax: B-cell lymphoma protein 2, *Bcl-2*: BCL2-Associated X Protein, mRNA: Messenger RNA, JAK2/STAT3: Janus kinase 2/signal transducer and activator of transcription 3, PDPN: Podoplanin, CLEC-2: C-type lectin-like receptor 2, MMP: Matrix metalloproteinase, PI3K: Phosphatidylinositol-3-kinase, TOPK: T-lymphokine-activated killer-cell-originated protein kinase, TxA2: Thromboxane A2, CDK4: Cyclin-dependent kinase 4, HepG2: Hepatoma G2, HIF-1α-FOXO3: Hypoxia-inducible factor 1-α-Forkhead Box O3, MEF2A: Myocyte enhancer factor 2A, JNK: Jun N-terminal kinase, ERK: extracellular regulated kinase, p-Akt: Phospho-protein kinase B, p-STAT3: Phospho-signal transducer and activator of transcription 3, AMP: Adenosine monophosphate, PPARα: Peroxisome proliferator activated receptor alpha, CPT-1α: Carnitine palmitoyltransferase -1α, PDK-4: Pyruvate dehydrogenase kinase 4, GLUT-4: Glucose transporter type 4, PI3K/Akt: PI3K and protein kinase B, LDH: Lactate dehydrogenase, CK-MB: Creatine kinase-myoglobin binding, GSH-PX: Glutathione peroxidase, Nrf2/HO-1: Nuclear factor erythroid 2-related factor transcription factor/Hemoxygenase 1, NF-κB: Nuclear factor kappa B, ICAM-1: Intercellular adhesion molecule 1, IL-6: Interleukin 6, EMPs: Endothelial microparticles, eNOS: Endothelial nitric oxide synthase, FGF2: Fibroblast growth factor-2, uPA: Urokinase-type plasminogen activator, CTGF: Connective tissue growth factor, TGF-β1: Transforming growth factor-β1, O-D: Ophiopogonin-D, *A. indica*: Anisomeles indica, *A. argyi*: *Artemisia argyi*, *C. morifolium*: *Chrysanthemum morifolium*, *G. glabra*: *Glycyrrhiza glabra*, *H. cordata*: *Houttuynia cordata*, *O. japonicus*: *Ophiopogon japonicus*, *P. frutescens*: *Perilla frutescens*, *P. grandiflorus*: *Platycodon grandiflorus*, MCF: Michigan cancer foundation

demonstrated that phytochemical substances extracted from *C. morifolium* Ramat flower extracts had a potent inhibitory effect on breast cancer cell lines (MCF-7 cells) [53]. According to gene ontology analysis, *Chrysanthemum*'s active compounds inhibit the growth of breast cancer through the regulation of the biological processes of their protein targets. 18-β-glycyrrhetic acid, glycycomarin, and glycyrrhizic acids present in *G. glabra* showed anti-cancer action in colon, liver, and lung cancer cell lines. The anti-cancer activity of 18-β-glycyrrhetic acid was evaluated against colorectal cancer cells LoVo, SW480, and SW620 [54]. The findings revealed that 18-β-glycyrrhetic acid might reduce PI3K and STAT3 signaling pathways as it regulates the apoptosis, invasion, and migration of colorectal cancer cells. Song *et al.* [55] reported that glycycomarin is highly effective against liver cancer in both cell culture and HepG2 xenograft models. The anti-cancer effect of glycycomarin is related to its capacity to directly inactivate T-lymphokine-activated killer-cell-originated protein kinase, resulting in the p53-dependent reduction of cell growth and induction of apoptosis. In an animal model of lung cancer, glycyrrhizin provided anti-cancer activity with low toxicity through inhibition of thromboxane A2 [56]. However, few studies investigated the effects of anti-proliferation, anti-invasion, anti-migration, and apoptosis induction by *H. cordata* extracts on different breast cancer cell types. For instance, the research by Subhawa *et al.* [57] showed that *H. cordata* Thunb extracts promoted breast cancer cell cycle arrest by

modulating the expressions of CDK4, cyclin D1, and p21 proteins. Moreover, by reducing matrix metalloproteinases release, the extracts reduced breast cancer cell migration and invasion. Furthermore, *H. cordata* Thunb induces apoptosis in human HepG2 hepatocellular cancer cells by activating the HIF-1A-FOXO3 and MEF2A signaling pathways [58]. Two important steroid glycosides, ophiopogonin B and D, isolated from *O. japonicus* exerted anti-cancer effects against human gastric cancer cells and breast carcinoma MCF-7 cells. Zhang *et al.* [59] reported that ophiopogonin B treatment resulted in increased phosphorylation levels of JNK1/2 and ERK1/2 in SGC-7901 cells. The anti-cancer mechanisms of ophiopogonin B may involve the JNK1/2 and ERK1/2 pathways. Another compound, ophiopogonin-D, reduced cell growth and promoted apoptosis in MCF-7 human breast cancer cells [60]. Ophiopogonin D-induced G₂/M cell cycle arrest was associated with downregulation of cyclin B1, and apoptosis was associated with the activation of caspase-8 and caspase-9. A study by Abd El-Hafeez *et al.* [61] showed a synergistic tumor-suppressive potency of *P. frutescens* and anti-cancer tyrosine kinase inhibitors on A549 cells, both *in vitro* and *in vivo*. The synergistic anti-tumor effect results from the activation of a two-stage cell cycle arrest at the G1 and G2/M phases. Another study showed that metabolites of *Platycodon grandiflorus*-induced cell apoptosis and inhibited cell proliferation by affecting the Akt and STAT3 signaling pathways [62]. Furthermore, a bioactive component of *Platycodon grandiflorum* called Platycodin D

causes cancer cell death associated with severe vacuolation through AMPK activation when cellular energy levels are low [63].

Cardiovascular protection

The JSHT ingredients and their phytochemical compounds demonstrate interventional activity in the process of cardiac remodeling and functional deterioration [Table 3]. In spontaneously hypertensive rats, *A. argyi* aqueous extract demonstrated antihypertensive activity, and a part of the mechanism appears to involve an increase in urine and electrolyte production [64]. With further research, *C. morifolium* Ramat showed similar antihypertensive activity [65]. The polyphenol-rich extract of *C. morifolium* Ramat ameliorated hypertension-induced cardiac hypertrophy in rats by lowering blood pressure and decreasing HIF-1 α expression, followed by regulation of PPAR-mediated PDK-4, GLUT-4, and CPT-1 α expressions. Furthermore, 18 β -Glycyrrhetic acid (extracted from *G. glabra*) demonstrated protective effects against myocardial infarction [7], indicating that it might have cardioprotective effects against acute myocardial infarction by suppression of inflammation, oxidative stress, and apoptosis via the PI3K/Akt pathway and reduction of Ca²⁺ concentration and cell contractility through L-type Ca²⁺ channels. A similar cardioprotective effect of *G. glabra* extract (GBE) was found in the DOX-induced cardiotoxicity mice model [66]. Mice treated with GBE were significantly protected from DOX-induced cardiotoxicity, as seen by lower creatine kinase isoenzyme and blood lactate dehydrogenase levels, improved heart morphology, and elevated glutathione levels and glutathione peroxidase activity. Similarly, 8-formyllophipogonanone B, a natural isoflavone in *O. japonicas*, protected against DOX-cardiotoxicity by reducing heme oxygenase-1-dependent myocardial inflammation and fibrosis [68]. Another compound, Platycodin D from *P. grandifloras*, considerably attenuates cardiac fibrosis [71]. A recent study revealed that the administration of *P. grandifloras* under conditions of spontaneous hypertension leads to a reduction in myocardial collagen content and fibrosis (downregulated the expression of fibroblast growth factor-2, urokinase-type plasminogen activator, matrix metalloproteinase-2, matrix metalloproteinase-2, and connective tissue growth factor and decreased transforming growth factor- β 1 expression). It is worth mentioning that some cardiovascular diseases, such as PM_{2.5}-mediated heart injury, have been linked to exposure to air pollution. Interestingly, by activating the Nrf2 transcription factor, 2-undecanone extracted from *H. cordata* can reduce the NF- κ B pathway and the inflammatory damage to mouse myocardium caused by exposure to PM_{2.5} [67]. In addition, inflammation of the blood vessels is critical to the development of cardiovascular diseases, including atherosclerosis. Paradee *et al.* showed that ethanol extracts of *P. frutescens* had a protective effect on endothelium against vascular inflammation by inhibiting intracellular adhesion molecule-1 expression and IL-6 production, adherence of leukocytes to endothelial cells, and endothelial microparticle generation [69]. Furthermore, another study

demonstrated that Perillaldehyde, one of the principal oil components in *P. frutescens*, inhibits atherosclerosis progression by endothelial nitric oxide synthase recovery and boosts tetrahydrobiopterin production [70].

Anti-diabetic activities

Globally, diabetes is one of the major chronic diseases, and approximately 10% of the world's population has a blood glucose metabolic abnormality characterized primarily by hyperglycemia. This condition is defined by either insufficient insulin secretion from β -cells of pancreatic islets, known as type 1 diabetes, or the inability of cells to react in response to secreted insulin, known as type 2 diabetes. A prolonged state of hyperglycemia causes increased ROS formation and dyslipidemia, resulting in extensive cellular damage and associated problems. Previous literature showed the potential effects of the ingredients of JSHT for diabetes management [Table 4]. The anti-diabetic effects of the flavonoids and phenolic compounds of *A. indica* (L.) Kuntze has been reported in experimental diabetic rats [72]. The results showed that ethyl acetate and hexane fraction of *A. indica* (L.) Kuntze improved fasting blood glucose, hemoglobin, and plasma insulin levels, and insulin sensitivity/resistance indicators such as quantitative insulin sensitivity check index, homeostasis model assessment of β -cell dysfunction, homeostasis model assessment of insulin resistance, and glucose tolerance. However, the aqueous extract of *A. argyi* exhibited marked anti-hyperglycemic effects in ALX-induced diabetic rats [73]. The aqueous extract showed marked improvements in serum insulin, total proteins, C peptide, and total hemoglobin, as well as decreases in glycosylated hemoglobin. Recently, Yamashita *et al.* reported the anti-hyperglycemic activity of *G. glabra* flavonoid oil (GFO) in diabetic KK-A^y mice [75], revealing that GFO improves hyperglycemia by targeting glucose transporter type-4 translocation in skeletal muscles through AMPK and Akt activation. Kumar *et al.* [76] also recorded the anti-hyperglycemic activity of *H. cordata* Thunb. extract in STZ-induced diabetic rats. The anti-diabetic effects of *H. cordata* are attributable to an increased expression of glucose transporter type-2 and GLUT-4 proteins, potentiating insulin secretion from pancreatic β -cells. Kim *et al.* [78] showed the anti-hyperglycemic effects of *P. frutescens* and proposed that *P. frutescens* can modulate the AMPK pathway and inhibit gluconeogenesis in the liver to prevent and treat type 2 diabetes. Besides, Zheng *et al.* found *Platycodon grandiflorum* aqueous-ethanol extract to have hypoglycemic effects without stimulating insulin secretion in STZ-induced diabetic ICR mice [79]. In obese diabetic KK-A^y mice, the hot water extract of *C. morifolium* Ramat showed hypoglycemic effects [74]. The mechanism involved suppression of chronic inflammation in adipose tissues, decreased adipocyte sizes, and increased peroxisome proliferator-activated receptor γ expression. Long-term hyperglycemia has been well documented as the primary cause of numerous diabetic complications, such as nephropathy and neuropathy. Diabetic nephropathy (DN) is a gradual consequence of diabetes, which may initially lead to nephrotic syndrome and subsequently evolve into chronic kidney disease and end-stage renal disease. Recently, Qiao *et al.* [77] showed the protective effect of ophiopogonin

Table 4: Anti-diabetic and anti-obesity effects of Jing Si herbal tea ingredients

Property	Sample/ingredients	Experimental model	Outcomes	Mode of action	Reference
Anti-diabetic	<i>A. indica</i> (L.) Kuntze	Animal model	Showed anti-diabetic activity in STZ-induced diabetic rats	Improved FBG, HbA1c, plasma insulin levels and HOMA-IR, HOMA- β and QUICKI	[72]
	<i>A. argyi</i>	Animal model	Showed anti-diabetic activity in ALX-induced diabetic rats	Marked improvements in serum insulin, total proteins, C peptide, and total hemoglobin, and decreases in HbA1c	[73]
	<i>C. morifolium</i> Ramat	Animal model	Exerted antidiabetic effects in obese diabetic KK-A ^y mice	Increased PPAR γ expression decreased the adipocyte sizes, suppressed chronic inflammation in adipose tissues, and elevated adiponectin levels	[74]
	<i>G. glabra</i>	Animal model	Exerted anti-hyperglycemic effect in diabetic KK-A ^y mice	Promoted GLUT4 translocation in skeletal muscle through AMPK and Akt activation	[75]
	<i>H. cordata</i>	Animal model	Exhibited anti-hyperglycemic effect in STZ-induced diabetic rats	Reversed the expression patterns of GLUT-2, GLUT-4, and caspase-3 levels	[76]
	<i>O. japonicus</i>	Animal model	Ophiopogonin D showed a protective effect against STZ-induced diabetic nephropathy	Increased serum albumin and creatinine clearance. Decreased serum creatinine, blood urea nitrogen, TGF- β 1, and kidney hypertrophy	[77]
	<i>P. frutescens</i>	Animal model	Exerted beneficial effects on hyperglycemia, dyslipidemia, glucose, and insulin intolerance	Modulated the AMPK pathway and inhibited gluconeogenesis in the liver	[78]
	<i>P. grandiflorus</i>	Animal model	Induced hypoglycemic effects without stimulating insulin secretion	Decreased blood glucose levels	[79]
Anti-obesity	<i>A. argyi</i>	Animal model	Improved dyslipidemia in HFD-induced obese rat	Reduced TG, TC, LDL-c, serum ALT, and AST levels	[80]
	<i>C. morifolium</i> Ramat	Animal model	Attenuated obesity-associated inflammation	Modulated the muscle AMPK-SIRT1 pathway	[81]
	<i>G. glabra</i>	Clinical study	Efficiently improve the lipid profile in overweight and obese subjects	Decreased TC, LDL-c levels, TC/HDL-c, LDL-c/HDL-c ratios, and log of TG/HDL-c	[82]
	<i>H. cordata</i>	Animal model	Showed anti-obesity effect and reduced dyslipidemia in HFD-induced obese rats	Decreased TC, TG, and LDL	[83]
	<i>O. japonicus</i>	Animal model	β -d-fructan prevented HFD-induced obesity and increased energy expenditure in mice	Ameliorated plasma lipid profiles, decreased leptin secretion, attenuated hepatic lipid accumulation, and increased the expressions of genes related to lipid and energy metabolism in the liver	[84]
	<i>P. frutescens</i>	In vitro and animal model	Showed anti-obesity effects in HFD-induced obese rats	Prevented body weight gain, improved serum lipids and hepatic lipids, and reduced epididymal fat	[85]
	<i>P. grandiflorus</i>	Animal model	Suppressed HFD-induced obesity in mice	Increased allobaculum and the production of SCFAs (butyrate and propionate) and other carbohydrate-related metabolites	[86]

FBG: Fasting blood glucose, HbA1c: Glycosylated hemoglobin, HOMA-IR: Homeostasis model assessment of insulin resistance, HOMA- β : Homeostasis model assessment of β -cell dysfunction, QUICKI: Quantitative insulin sensitivity check index, STZ: Streptozotocin, ALX: Alloxan, PPAR γ : Peroxisome proliferator-activated receptor- γ , GLUT: Glucose transporter type, AMPK: 5-Prime-Amp-activated protein kinase, Akt: Protein kinase B, TGF- β 1: Transforming growth factor- β 1, HFD: High-fat diet, TG: Triglycerides, TC: Total cholesterol, HDL-C: High-density lipoprotein cholesterol, LDL-C: Low-density lipoprotein-cholesterol, ALT: Alanine aminotransferase, AST: Aspartate transaminase, AMPK-SIRT1: AMPK-silent information regulator T1, SCFAs: Short-chain fatty acids, *A. indica*: *Anisomeles indica*, *A. argyi*: *Artemisia argyi*, *C. morifolium*: *Chrysanthemum morifolium*, *G. glabra*: *Glycyrrhiza glabra*, *H. cordata*: *Houttuynia cordata*, *O. japonicus*: *Ophiopogon japonicus*, *P. frutescens*: *Perilla frutescens*, *P. grandiflorus*: *Platycodon grandiflorus*

D extracted from *O. japonicas* against renal damage in an STZ-induced DN rat model. Ophiopogonin D treatment decreased TGF- β 1, serum creatinine, kidney hypertrophy, and blood urea nitrogen. Furthermore, serum albumin and creatinine clearance were increased by ophiopogonin D treatment in DN rats. Thus, the literature shows that *O. japonicas* could alleviate the development and progression of DN.

Anti-obesity activities

Over the past few decades, obesity has become a major health issue worldwide. Metabolic syndrome symptoms linked to obesity include hypertension, glucose intolerance,

insulin resistance, and dyslipidemia. Obesity in humans can be prevented and managed through various approaches, and efforts have been made to identify new materials since obesity cannot be entirely cured or prevented, and the vast majority of anti-obesity drugs can lead to adverse effects. The ingredients of JSHT are recognized to have an anti-obesity impact and are thus used to treat obesity [Table 4]. Investigating the anti-obesity activity of *A. argyi* in high-fat diet (HFD)-induced obese rats, a study found that *A. argyi* extracts significantly lower the levels of TG, TC, LDL-c, serum ALT, and AST levels [80]. These results imply that

by increasing lipid metabolism, *A. argyi* extracts may aid in weight loss and dyslipidemia treatment. A similar study was also conducted on *H. cordata* extract by Thunb *et al.* [83] < which showed that the extract decreased the levels of TC, TG, and LDL in HFD-induced obese rats. Correspondingly, rodents treated with *P. frutescens* also showed reduced obesity through prevention of body weight gain, improvement of hepatic lipids and serum lipids, and reduction of epididymal fat [85]. Furthermore, Mirtaheri *et al.* [82] conducted a double-blind, randomized, controlled clinical trial to investigate the effects of GBE on the lipid profile and atherogenic indices in 64 overweight and obese subjects. Their findings revealed that after 8 weeks, GBE decreased the levels of TC, LDL-c levels, TC/HDL-c, LDL-c/HDL-c ratios, and log of TG/HDL-c. Decreased energy expenditure and chronically positive energy balance contribute to the prevalence of obesity and associated metabolic dysfunctions such as dyslipidemia, hepatic fat accumulation, inflammation, and muscle mitochondrial defects. Wang *et al.* asserted that β -D-fructan, a water-soluble polysaccharide extracted from *Ophiopogon japonicus*, is a promising candidate drug for the treatment of obesity-related metabolic diseases due to its ameliorative effect on lipid profile, attenuation of hepatic lipid accumulation, and upregulation of 12 genes involved in lipid and energy metabolism in the liver [84]. Lee *et al.* [81] examined the effects of *C. morifolium* Ramat flower extract on obesity-induced inflammation. By regulating the AMPK-SIRT1 pathway in muscles, the extract reduces the inflammation associated with obesity. Recently, gut microbiota has been a potential target for anti-obesity dietary interventions. A diet rich in bioactive compounds, including polyphenols and dietary fiber, can regulate gut flora and promote healthy weight maintenance. A study by Ke *et al.* [86] showed that *Platycodon grandiflorus* altered the gut metabolome and gut microbiota in HFD-fed mice; it primarily increased allobaculum and the production of SCFAs (butyrate and propionate) and other carbohydrate-related metabolites.

Hepatoprotective properties

Chronic liver disease, such as nonalcoholic steatohepatitis, nonalcoholic liver disease, liver fibrosis, cirrhosis, and many other conditions, can be caused by a wide range of chemicals, including drugs, dietary components, alcohols, and other environmental pollutants. Excessive alcohol consumption can cause liver dysfunction and alcoholic liver disease, resulting in liver damage. Researchers have proven that AST, ALT, ALP, and bilirubin can be expressed in the liver and that their abnormally high levels can cause damage and necrosis in liver tissues [Table 5]. A recent study showed that the aqueous and ethanolic extracts of *A. indica* (L.) Kuntze decreased ALT, AST, and ALP levels in STZ-induced diabetic rat models [87]. Liver cells are sensitive to oxidative stress, and it was reported that *A. argyi* extract exhibited protective activity against chemically and immunologically induced acute liver injuries in a CCl_4 -induced liver injury mouse model of [88]. According to the findings, *A. argyi* may help prevent oxidative liver damage, maintain intracellular levels of antioxidant enzymes, reduce levels of ALT, AST,

ALP, TNF- α , IL-1, and MDA, and improve SOD and GPx activities. In an APAP-induced liver injury mouse model, the ethyl acetate extract of *C. morifolium* Rama, which protects against liver injury, showed significant anti-oxidant activity [89]. Mechanistically, *C. morifolium* Rama might reduce apoptosis through PI3K–Akt pathway and suppress excessive oxidative stress through the GSK3 β –Nrf2 pathway. In a laboratory experiment, licorice total flavonoids (LTF) were isolated from *G. glabra*, and three different types of liver damage mice models were used to study its hepatoprotective effects, including tetrachloromethane, Chinese liquor, and a high-fat emulsion [90]. Their findings suggested that LTF can reduce hepatic injury and repair liver tissue by reducing the levels of AST, ALT, ALP, TG, cholesterol, hepatic MDA, and increasing hepatic SOD and GPx of hepatic injury mice. Furthermore, a recent report from Kang and Koppula [91] claimed the hepatoprotective effects of *H. cordata* ethyl acetate extract in CCl_4 -treated mice due to its significant reduction in the levels of increased serum and regulation of the altered levels of serum cholesterol. Recently, there has been a surge in interest in the polysaccharides derived from *O. japonicas* roots, which exhibit antioxidant properties and protect the liver from the detrimental effects of diabetes. Chen *et al.* [92] showed that OJP1, a polysaccharide isolated from the roots of *Ophiopogon japonica*, significantly reduced the levels of TG, TC, LDL-C, HDL-C, and MDA while increasing the activities of SOD and GPx in diabetic rats. Furthermore, the hepatoprotective effects of rosmarinic and caffeic acid, two primary compounds of *P. frutescens*, against *t*-BHP-induced oxidative liver damage were reported by Yang *et al.* [93]. Both compounds reduced MDA, ALT, and AST levels and increased GSH, GPx, and SOD activity. Moreover, the attenuative effects of *Platycodon grandiflorus* on hepatic insulin resistance and oxidative stress in HFD-induced nonalcoholic fatty liver diseases were investigated [94]. By modulating fatty acid synthase expression and phosphorylation of acetyl-CoA carboxylase, *Platycodon grandiflorus* significantly decreased HFD-induced liver damage, hyperlipidemia, and hepatic steatosis. *Platycodon grandiflorus* also improves insulin sensitivity by regulating the PI3K/Akt/GSK3 β signaling pathway and alleviating oxidative stress by increasing glutathione content and antioxidant activities.

Anti-inflammatory activities

The anti-inflammatory activity of JSHT ingredients has been tested both *in vitro* and *in vivo* [Table 5]. Nasrin *et al.* [16] investigated the *in vitro* anti-inflammatory potential of a methanol extract of *A. indica* (L.) Kuntze (MeOH-AI), revealing that MeOH-AI significantly inhibited protein denaturation and hemolysis induced by heat and hypotonic solution. Computer-aided analyses and molecular docking indicated anti-inflammatory activity through binding with COX-2. Moreover, inflammatory bowel diseases (IBD), like ulcerative colitis, have become more prevalent globally. Shin *et al.* [95] reported that in a dextran sulfate sodium-induced colitis mouse model, *A. argyi* ethanol extracts mediated IBD symptoms and enhanced immunomodulatory responses in lymphoid tissues through the activation of Nrf2 and

Table 5: Hepatoprotective and anti-inflammatory effects of Jing Si herbal tea ingredients

Property	Sample/ ingredients	Experimental model	Outcomes	Mode of action	Reference
Hepatoprotective	<i>A. indica</i> (L.) Kuntze	Animal model	Aqueous and ethanolic extracts showed hepatoprotective effects STZ-induced diabetic rat models	Reduced the levels of ALT, AST, ALP	[87]
	<i>A. argyi</i>	Animal model	Protected against oxidative liver injury	Reduced the levels of ALT, AST, ALP, TNF- α , IL-1, and MDA. Restored the activities of SOD and GPx	[88]
	<i>C. morifolium</i> Ramat	Animal model	Showed protective effect from APAP-induced liver injury	Reduced the levels of AST, ALT, and ROS. Upregulated SOD and GSH. Increased levels of p-AMPK, p-GSK3 β , HO-1, and NQO1, and promoted Nrf2	[89]
	<i>G. glabra</i>	Animal model	Reduce hepatic injury in mice	Reduced ALT, AST, and ALP, hepatic MDA, TG, and cholesterol levels. Increased SOD and GPx activities	[90]
	<i>H. cordata</i>	Animal model	Exhibited significant hepatoprotective property in carbon tetrachloride-induced hepatotoxicity in mice	Reduced the elevated serum levels and regulated the altered levels of serum cholesterol	[91]
	<i>O. japonicus</i>	Animal model	Showed liver-protective effects from the injurious effects of diabetes	Reduced the levels of TG, TC, LDL-C, HDL-C, and MDA. Increased the activities of SOD and GPx in the serum	[92]
	<i>P. frutescens</i>	Animal model	Showed hepatoprotective effect against oxidative liver injury	Reduced the levels of AST, ALT, and MDA and increased the activities of GSH, GPx, and SOD	[93]
	<i>P. grandiflorus</i>	Animal model	Exerted profound effects on hepatic insulin sensitivity and oxidative stress	Regulated phosphorylation of acetyl-CoA carboxylase, expression of fatty acid synthase, and PI3K/Akt/GSK3 β signaling pathway	[94]
	<i>A. indica</i> (L.) Kuntze	<i>In vitro</i>	Anti-inflammatory studies produced remarkable responses	Inhibited protein denaturation and hemolysis induced by heat and hypotonic solution	[16]
Anti-inflammatory	<i>A. argyi</i>	<i>In vitro</i> and animal model	Mediated IBD symptoms and enhanced immunomodulatory responses in lymphoid tissues	Activated Nrf2, and HO-1, inhibited NF- κ B, and upregulated IL-10	[95]
	<i>C. morifolium</i> Ramat	<i>In vitro</i>	Inhibited the production of inflammatory mediators	Inhibited TLR4 signaling	[96]
	<i>G. glabra</i>	Animal model	Diammonium Glycyrrhizinate reduced inflammatory injury	Suppressed NF- κ B, TNF- α , and ICAM-1 in colonic mucosa	[97]
	<i>H. cordata</i>	<i>In vitro</i> and animal model	Showed anti-inflammation potential	Reduced NO production and suppressed the expression of PGE2, iNOS, IL-1 β , TNF- α , and IL-6	[98]
	<i>O. japonicus</i>	<i>In vitro</i>	Showed anti-inflammation potential	Inhibited the phosphorylation of ERK1/2 and JNK in MAPK signaling pathways	[99]
	<i>P. frutescens</i>	Animal model	Protected against dextran sulfate sodium-induced murine colitis	Inhibited the activation of NF- κ B and STAT3 and elevated the accumulation of Nrf2 and HO-1	[100]
	<i>P. grandiflorus</i>	<i>In vitro</i>	Showed anti-inflammation potential	Suppressed NF- κ B and activated HO-1	[101]

STZ: Streptozotocin, ALT: Alanine aminotransferase, AST: Aspartate transaminase, ALP: Alkaline phosphatase, TNF- α : Tumor necrosis factor-alpha, IL: Interleukin, MDA: Malondialdehyde, SOD: Superoxide dismutase, GPx: Glutathione peroxidase, ROS: Reactive oxygen species, GSH: Glutathione, p-AMPK: Phosphorylated - 5-prime-amp-activated protein kinase, p-GSK3 β : Phosphorylated-glycogen synthase kinase 3 Beta, HO-1: Hemoxygenase 1, NQO1: NAD (P) H quinone dehydrogenase 1, TG: Triglyceride, TC: Total cholesterol, LDL-C: Low-density lipoprotein-cholesterol, HDL-C: High-density lipoprotein cholesterol, PI3K/Akt/GSK3 β : Phosphatidylinositol-3-kinase and protein Kinase B/glycogen synthase kinase 3 Beta, IBD: Inflammatory bowel diseases, Nrf2: Nuclear factor erythroid 2-related factor 2, TLR4: Toll-like receptor 4, ICAM-1: Intercellular adhesion molecule 1, NO: Nitric oxide, PGE2: Prostaglandin E2, iNOS: Inducible nitric oxide synthase, IL-1 β : Interleukin 1 beta, ERK1/2: Extracellular-regulated kinase 1/2, IL-6: Interleukin 6, JNK: Jun N-terminal Kinase, MAPK: Mitogen-activated protein kinase, STAT3: Signal transducer and activator of transcription 3, NF- κ B: Nuclear factor kappa B, *A. indica*: *Anisomeles indica*, *A. argyi*: *Artemisia argyi*, *C. morifolium*: *Chrysanthemum morifolium*, *G. glabra*: *Glycyrrhiza glabra*, *H. cordata*: *Houttuynia cordata*, *O. japonicus*: *Ophiopogon japonicus*, *P. frutescens*: *Perilla frutescens*, *P. grandiflorus*: *Platycodon grandiflorus*, APAP: Acetaminophen

HO-1, inhibition of NF- κ B, and upregulation of IL-10. Similar result were also reported by Park *et al.* [100] for *P. frutescens* extracts. *P. frutescens* treatment prevented the activation of NF- κ B and STAT3 produced by dextran sulfate sodium, although it increased the accumulation of Nrf2 and HO-1 in the colon. Correspondingly, the anti-inflammatory mechanism of *P. grandifloras* ethanol extract was associated with the inhibition of NF- κ B translocation

and the activation of HO-1 production [101]. Further, to explore a possible anti-inflammatory effect of diammonium glycyrrhizinate (extracted from *G. glabra*) in ulcerative colitis, the suppressive expression of TNF- α , ICAM-1, and NF- κ B in colonic mucosa was detected by immunohistochemistry in a rat model [97]. Furthermore, in LPS-stimulated RAW264.7 cells, the methanolic and aqueous extracts of *H. cordata* fermentation product inhibited the generation of PGE2, NO,

and inflammatory cytokines (TNF- α , IL-1, IL-6) [98]. Later, a carrageenan-induced rat paw edema model also confirmed the anti-inflammatory activities of fermented *H. cordata*. Zhao *et al.* isolated thirteen homoisoflavonoids compounds from *O. japonicas* [99]. Among these isolated compounds, 4'-*O*-demethylophiopogonanone E exhibited anti-inflammatory effect through suppressing the phosphorylation of JNK and ERK1/2 in MAPK signaling pathways, hence reducing the generation of pro-inflammatory cytokines and NO. Recently, a new bisepoxylignan called dendranlignan A, isolated from *C. morifolium* Ramat flowers, inhibited LPS-induced inflammation in H9c2 cardiomyocytes via the TLR4 pathway [96].

Kidney protective activities

Chronic kidney disease is typically caused by oxidative damage, inflammation, and other factors that interfere with the water and salt metabolism in the body [Table 6]. Antioxidative compounds can protect against contrast agent-induced cytotoxicity in renal proximal tubular cells and preserve renal function during the pathogenesis of an acute kidney injury. For instance, Kim *et al.* isolated 14 phenolic compounds from *A. argyi*. Their findings indicated that the anti-oxidant and antiapoptotic effects of methyl caffeate, an isolated phenolic compound from the *A. argyi*, protect LLC-PK1 cells from iodixanol-induced cell death. This action is attributed to MAPK (P38, JNK, and ERK) inhibition as well as KIM-1 and caspase-3 activities [102]. Xia *et al.* [103] found that total flavonoids from *C. morifolium* Ramat exerted kidney protective effects against lead-induced oxidative damage in mice by increasing CAT and GSH-Px activity and decreasing MDA concentration. Furthermore, ophiopogonin A, an active compound of *O. japonicas*, alleviated hemorrhagic shock-induced renal injury via the p-ERK/ERK signaling pathway [106]. Inflammation also contributes significantly to kidney disease. Zhang *et al.* [108] showed that saponins extracted from the roots of *Platycodon grandiflorum* protect mice kidneys from cisplatin-induced damage by modulating PI3K/Akt/apoptosis signaling pathways and suppressing NF- κ B activation. Notably, most kidney damage or nephrotoxicity occurs as a result of drug use and is much more common in hospitalized patients. *G. glabra*, another important component of JSHT, demonstrated an ameliorative impact against nephrotoxicity induced by gentamicin in mice [104]. Gentamicin increased tissue levels of Bax, and Cox-2 and decreased expression of Nrf-2, and HO-1, but these levels recovered to normal in the *G. glabra* -treated group. Interestingly, rats treated with *H. cordata* exhibited a considerable improvement in renal function, most likely due to lowered biochemical indices and oxidative stress parameters (decreased levels of MDA, BUN, and creatinine and increased levels of CAT, SOD, and GSH) associated with gentamicin-sulfate-induced nephrotoxicity [105]. Multiple studies support the notion that renal ROS mediates kidney damage in patients with diabetes. *P. frutescens* sprout extract reportedly prevents high-glucose-induced renal mesangial cell dysfunction by modulating AMPK and NADPH oxidase signaling [107].

Neuroprotective activities

An alarming rise has been observed in the prevalence of neurodegenerative diseases. Despite advances in the treatment of Alzheimer's, Parkinson's, and Huntington's diseases, the pathophysiology of these conditions is not yet entirely understood. The provision of neuroprotective agents may delay the onset and reduce the symptoms of neurodegenerative diseases by decreasing inflammation, apoptosis, immune dysfunction, and oxidative stress. Numerous studies have established the potential neuroprotective properties of JSHT's ingredients [Table 6].

One experimental study suggested that Alzheimer's disease (AD) may benefit from the ethyl acetate extract of *A. indica* (L.) Kuntze, which may have anti-oxidant and acetylcholinesterase (AChE) inhibitory properties due to its phenolic components [109]. AD results from chronic inflammation of certain parts of the brain and its development could be attributed to oxygen-free radicals in elderly individuals. JSHT's ingredients may provide protective effects through their antioxidant properties, resulting in a reduced risk of brain damage and improved neurological function. It has been suggested that a combination of anti-inflammatory and antioxidant activities with neuroprotective effects might result in memory improvement. In a test of animal behavior, 3,5-dicaffeoylquinic acid (3,5-diCQA)—the primary phenolic ingredient in *A. argyi*—dramatically ameliorated learning and memory deficits by boosting the levels of AChE and SOD while decreasing the levels of AChE, MDA, and GSH [111]. In addition, 3,5-diCQA may prevent neuronal death by protecting mitochondrial functions and inhibiting apoptotic signaling molecules such as p-Akt, Bax, and p-tau (Ser 404). Similarly, a study showed that *H. cordata* water extracts improved symptoms of cognitive impairment in cholinergic dysfunction AD's disease models due to cholinergic dysfunction and inhibited cholinergic antagonists and tauopathies [120]. Furthermore, glycyrrhizic acid, another important phytochemical compound isolated from *G. glabra*, significantly improved copolamine-induced cognitive impairment in a mouse model [117]. Administration of *G. glabra* increased phosphorylation of ERK and JNK proteins, SOD, and CAT enzyme activities, and decreased the activity of AChE. A β is well established as one of the initiators that causes the progression of AD by accumulation and aggregation, which can be caused by A β overproduction or clearance failures. Wang *et al.* [114] isolated luteolin, a flavonoid compound of *C. morifolium* Ramat and found that it exhibited a neuroprotective effect in an STZ-induced AD rat model by dispersing A β plaques and inhibiting free radical products. Furthermore, the ethyl acetate fraction of the methanol extract of *Ophiopogon japonicas* displayed significant protective effects against A β -induced cytotoxicity to PC12 cells and was a potent inhibitor of β -secretase activity, resulting in decreased A β production [121]. Similarly, in 5XFAD mice, the ethanol extract of leaves of *P. frutescens* inhibited and disassembled A β aggregation, lowering lipid peroxidation and bad cholesterol [123]. However, Nam *et al.* [125] suggested

Table 6: Nephron- and neuroprotective effects of Jing Si herbal tea ingredients

Property	Sample/ ingredients	Experimental model	Outcomes	Mode of action	Reference
Nephroprotective	<i>A. argyi</i>	<i>In vitro</i>	Protected kidney epithelium cells against apoptotic damage caused by iodixanol	Inhibited the activities of MAPKs (JNK, P38 and ERK), caspase-3, and KIM-1	[102]
	<i>C. morifolium</i> Ramat	Animal model	Showed kidney protective effects against lead-induced oxidative damage in mice	Increased the activity of GSH-Px, and CAT and decreased MDA concentration	[103]
	<i>G. glabra</i>	Animal model	Showed ameliorative impact against nephrotoxicity induced by gentamicin	Increased the levels of SOD and GSH-Px, promoted the expression of Nrf2 and COX-2, and inhibited IL-1 β and IL-6 production	[104]
	<i>H. cordata</i>	Animal model	Showed protective effects against nephrotoxicity and oxidative stress	Decreased the levels of creatinine, BUN, and MDA. Increased the levels of GSH, SOD, and CAT	[105]
	<i>O. japonicus</i>	<i>In vitro</i> and animal model	Alleviated hemorrhagic shock-induced renal injury	p-ERK/ERK signaling pathway	[106]
	<i>P. frutescens</i>	<i>In vitro</i>	Protected renal mesangial cell dysfunction against high glucose	Modulated AMPK and NADPH oxidase signaling	[107]
	<i>P. grandiflorus</i>	Animal model	Exerted kidney protection effects against cisplatin-induced kidney injury	Inhibiting the activation of NF- κ B and regulated PI3K/Akt/apoptosis signaling pathways	[108]
Neuroprotective	<i>A. indica</i> (L.) Kuntze	<i>In vitro</i>	Showed potential effect on AD treatment	Inhibited cholinesterase activities	[109]
	-	Animal model	Ovatodioidide prevented rats from brain I/R damage	Hampered neuronal apoptosis and microglial inflammation via the SIRT1-NF- κ B pathway	[110]
	<i>A. argyi</i>	Animal model	Ameliorated the cognitive impairment induced by TMT	Prevent neuronal apoptosis by protecting mitochondrial activities and repressing apoptotic signaling molecules such as p-Akt, BAX, and p-tau (Ser 404)	[111]
	-	<i>In vitro</i>	Showed protective effects against PD	Reduced ROS, cell apoptosis, and MDA levels. Increased GSH and SOD levels	[112]
	-	<i>In vitro</i>	Eupatilin showed neuroprotective effects against focal cerebral ischemia	Reducing IKK α / β phosphorylation, I κ B α phosphorylation, and I κ B α degradation	[113]
	<i>C. morifolium</i> Ramat	Animal model	Luteolin showed a protective effect on learning defects and hippocampal structures in AD	Inhibited free radical products and dispersing A β plaques	[114]
	-	Animal model	Showed protective effects against PD	Increased the levels of SOD and GSH-Px and decreased MDA levels	[115]
	-	Clinical trial	Showed neuroprotective effect on ischemic stroke patients	Increased the levels of BDNF, SOD, and TAC and decreased the levels of the NIHSS, serum NSE, S100, and MDA	[116]
	<i>G. glabra</i>	Animal model	Showed neuroprotective effect on cognitive function	Increased phosphorylation of ERK and JNK proteins. Decreased activity of AChE and increased activity of SOD and CAT enzymes	[117]
	-	Clinical trial	Improved the symptoms in PD patients	Improved total UPDRS scores, daily activities and tremors along with motor test and rigidity scores	[118]
	-	Clinical trial	Improved the symptoms in patients with acute ischemic stroke	Declined in NIHSS scores and MRS	[119]
	<i>H. cordata</i>	Animal model	Improved cognitive deficits in cholinergic dysfunction AD's disease-like models	Inhibited cholinesterase activities	[120]
	<i>O. japonicus</i>	<i>In vitro</i>	Exhibited a neuroprotective effect on A β toxicity	Inhibited β -secretase activity and decreased MDA content	[121]
	-	Animal model	Methylophipogonanone A showed therapeutic potential against cerebral I/R injury	Regulated the expression of MMP-9 and tight junction proteins	[122]
	<i>P. frutescens</i>	Animal model	Inhibited A β aggregates induce memory deficit	Inhibited and disassembled A β aggregation, lowered the lipid peroxidation and bad cholesterol	[123]

Contd...

Table 6: Contd...

Property	Sample/ ingredients	Experimental model	Outcomes	Mode of action	Reference
-		Animal model	Counteract VaD-associated structural and functional deterioration	Attenuated proinflammatory mediators, including NO, TNF- α , and IL-6, downregulated the upstream MAPKs/NF κ B/iNOS pathway	[124]
	<i>P. grandiflorus</i>	Animal model	Improved cognitive deficits	Decreased A β -related pathology and ameliorated A β -induced cognitive impairment	[125]
-		Animal model	Improved learning and memory by enhancing synaptogenesis	Activated the MAPK/ERK signaling pathway	[126]

MAPKs: Mitogen-activated protein kinases, KIM-1: Kidney injury marker-1, JNK: Jun N-terminal kinase, ERK: extracellular regulated kinase, GSH-Px: Glutathione peroxidase, CAT: Catalase, MDA: Malondialdehyde, SOD: Superoxide dismutase, Nrf2: Nuclear factor erythroid 2-related factor 2, COX-2: Cyclooxygenase-2, IL-1 β : Interleukin 1 beta, IL-6: Interleukin 6, BUN: Blood urea nitrogen, AMPK: Adenosine monophosphate-activated kinase, NADPH: Nicotinamide adenine dinucleotide phosphate hydrogen, NF- κ B: Nuclear transcription factor-kappa B, AD's: Alzheimer's disease, SIRT1: Sirtuin-1, TMT: Trimethyltin, BAX: B-cell lymphoma protein 2, PD: Parkinson's disease, ROS: Reactive oxygen species, IKK α / β : Inhibitory kappa B kinase α , A β : Amyloid-beta, BDNF: Brain-derived neurotrophic factor, TAC: Total antioxidant capacity, NIHSS: National Institutes of Health Stroke Scale, NSE: Neuron-specific enolase, AChE: Acetylcholinesterase, UPDRS: Unified Parkinson's disease rating scale, MRS: Modified Rankin scale, MMP-9: Matrix metalloproteinase-9, NO: Nitric oxide, TNF- α : Tumor necrosis factor-alpha, iNOS: Inducible nitric oxide synthase, PI3K/Akt: Phosphatidylinositol-3-kinase and protein kinase B, I κ B α : Nuclear factor of kappa light polypeptide gene enhancer in B-cells inhibitor, alpha, *A. indica*: *Anisomeles indica*, *A. argyi*: *Artemisia argyi*, *C. morifolium*: *Chrysanthemum morifolium*, *G. glabra*: *Glycyrrhiza glabra*, *H. cordata*: *Houttuynia cordata*, *O. japonicus*: *Ophiopogon japonicus*, *P. frutescens*: *Perilla frutescens*, *P. grandiflorus*: *Platycodon grandiflorus*

that as a potential therapeutic agent for AD, *Platycodon grandiflorus* root extract may be able to decrease A β -related pathology and ameliorate A β -induced cognitive impairment. Hippocampal synaptogenesis is a key process for cognitive function, which involves the formation of synapses between neurons in the hippocampus and adult hippocampal neurogenesis. Kim *et al.* [126] found that the root extract of *Platycodon grandiflorus* improved learning and memory by increasing synaptogenesis via activation of the ERK1/2 signaling pathway in the hippocampus of mice. Another neurodegenerative disorder, vascular dementia (VaD) is defined by time-dependent memory loss and is essentially accompanied by neuroinflammation. In a recent study, polyphenol-rich PLE exerted therapeutic efficacy against VaD via decreasing proinflammatory mediators, including NO, TNF- α , and IL-6, and downregulating the upstream MAPKs/NF- κ B/iNOS pathway [124].

The etiology of Parkinson's disease (PD), the second most prevalent age-related neurodegenerative disease, remains to be clarified. Inflammation and oxidative stress are crucial factors in the development and progression of PD. The ingredients of JSHT have been shown to protect cells from oxidative stress, inflammation, apoptosis, and mitochondrial dysfunction. One study evaluated the neuroprotective effects of *A. argyi* against PD on 6-hydroxydopamine-induced toxicity in SH-SY5Y cells [112], revealing that the ethanolic extracts of *A. argyi* significantly reduced ROS, cell apoptosis, and MDA levels, as well as increased GSH and SOD levels. Similar results were also recorded for *C. morifolium* Ramat extracts against 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine-induced PD in C57BL/6 mice [115]. In a clinical trial, the ingestion of licorice root (*G. glabra*) could alleviate the symptoms of PD without causing major adverse effects [118]. Four months following licorice consumption, total Unified PD Rating Scale scores, daily activities, tremor, motor test, and rigidity scores improved significantly.

Cerebral ischemia—including stroke—is the leading cause of disability and death worldwide. It causes irreversible damage to the brain and loss of neural function. Ischemia results in an increase in ROS, mitochondrial dysfunction, and neuronal death as a result of a lack of oxygen, impaired blood flow, and calcium accumulation. Hu *et al.* [110] demonstrated the ability of ovatodiolide (extracted from *A. indica* [L.] Kuntze) to attenuate neurological apoptosis and deficits following cerebral ischemia in a dose-dependent manner. Mechanistically, OVA prevented brain I/R damage in rats by inhibiting neuronal apoptosis and microglial inflammation through the SIRT1-NF- κ B pathway. Another compound, eupatilin—a substance isolated from *A. argyi*—was found to have neuroprotective benefits against focal cerebral ischemia by reducing NF- κ B signaling activity in the ischemic brain by decreasing IKK α / β phosphorylation, I κ B α phosphorylation, and I κ B α degradation [113]. Furthermore, a homoisoflavonoid compound of *O. japonicas* known as methylphopogonone A, showed therapeutic potential for the treatment of cerebral I/R injuries in a rat model due to its ability to mitigate blood–brain barrier damage via modulation of tight junction proteins and MMP-9 expression [122]. In the treatment of ischemic stroke, reducing oxidative stress and increasing neuroprotection can play a critical role. In a clinical trial, Zhu *et al.* [116] evaluated the protective effect of *C. morifolium* Ramat extract on patients with ischemic stroke (60 patients). They found that compared to the levels on the 2nd day, the levels of TAC, BDNF, and SOD were increased, and the MDA, serum NSE, NIHSS, and S100 levels were decreased in the *C. morifolium* Ramat group on the 4th day. In another clinical trial, Ravanfar *et al.* [119] evaluated the efficacy of the whole extract of licorice (extracted from *G. glabra*) in the neurological improvement of 75 patients after acute ischemic stroke. Results demonstrated that patients who ingested 450 mg and 900 mg of licorice extracts after 24 h of an ischemic stroke

had significantly better neurologic improvement (based on NIHSS scores) after 3 months.

TOXICITY OF THE INGREDIENTS OF JSHT

Despite well-established application, products derived from any herb require not only efficacy but also safety evidence. Several investigations on the toxicological effects of JSHT ingredients have been published, and surprisingly, most studies have shown weak potential toxicity. Basappa *et al.* [127] conducted a study to evaluate the toxicity of *A. indica* (L.) Kuntze leaf flavonoid fraction (ALFF) in mice, human lymphocytes, and human cancer cell lines. Under the experimental conditions, acute and subacute exposures to ALFF are nontoxic to mice, with the maximum tolerated dose exceeding 5000 mg/kg body weight. ALFF reportedly does not influence the hematological system, is not clastogenic or mutagenic, and inhibits the multiplication of cancer cells. Studies on the acute toxicity of hexane leaf extract of the *Artemisia* species were carried out by Ogbole *et al.* [128] on rats administered doses of 1000, 2000, and 2500 mg/kg intraperitoneally. They monitored and observed various parameters, such as the rats' decreased intake of food and fluids, in contrast to their increased output of urine and feces. Moreover, few significant changes were observed in the hematological system, and the extract was shown to have a lethal dose (LD50) of 2750 mg/kg body weight. Overall, studies showed low toxicity in short-term treatment. In a long-term toxicity study, Li *et al.* [129] treated the rats with *C. morifolium* Ramat daily by oral gavage at dose levels of 320, 640, and 1280 mg/kg body weight for 26 consecutive weeks followed by a 4 weeks recovery period. The data demonstrated no toxicological changes in food, water consumption, body weight, microscopic histopathologic examination, hematologic examination, organ weight, and blood biochemical examination were found in any treatment group. According to our assessments of the scientific literature, *G. glabra* is not a prominent teratogen and exhibits mild mutagenicity, genotoxicity, carcinogenicity, and developmental toxicity. Based on the scientific evidence, Nazari *et al.* [130] claimed that *G. glabra* is moderately toxic. Having said that, they also mentioned that since oral dispensation reduces absorption and first-pass metabolism of *G. glabra*, there is a lower risk of toxicity after oral administration than after intravenous or IP administration. However, the use of *G. glabra* and its components is contraindicated in pregnancy and neonates. As an edible plant, *H. cordata*'s potential toxicity is typically overlooked. There are no reports of its toxicity in long-term consumption as a vegetable or for medicinal use alone in certain regions of Asia, and the National Health Commission of China listed it in 2013 as one of the plants that can be used as both food and medicine, suggesting that *H. cordata* is relatively safe for oral administration in humans [20]. Although *O. japonicas* is well-known as both food and a therapeutic herb, the toxicity and safety assessments for this plant remain lacking. Few investigations on the toxicity of *O. japonicas* decoction have been conducted. Zhang *et al.* [131] evaluated the possible development of *O. japonicas* -induced toxicity in Sprague Dawley (SD) rats, and found that *O. japonicas* decoction (26.9 g/kg) had no significant effects on maternal body weight, fetal weight, viability, or fetal malformations in SD rats. The results showed that *O. japonicas* had no harmful effects on

the fetuses or on the pregnant female SD rats that were treated. Interestingly, there are very few studies that have reported on the toxicological aspects of materials originating from *P. frutescens*. The inhalation of smoke from roasted perilla seeds caused occupational asthma via an IgE-mediated mechanism. In addition, a single instance of anaphylaxis due to perilla seed has been documented [22]. Regarding the *P. grandifloras*, Lee *et al.* [132] reported that at doses up to 2000 mg/kg, playcodin D did not cause treatment-related mortality, significant changes in body weight, substantive histopathological changes, or abnormal clinical signs in 14 principal organs in male and female mice.

CONCLUSIONS AND FUTURE PERSPECTIVES

The eight ingredients of JSHT are potentially rich sources of phytochemicals that may help reduce disease risk conditions and, therefore, manage several noncommunicable diseases. Based on current evidence, this review summarized the therapeutic activities of JSHT's individual ingredients. It is clear from the reported evidence that the pharmacological activities of the eight ingredients in JSHT strengthen the promotion of JSHT as a health-care product. Thus, the review will provide a new foundation for consumers and the scientific community to better understand the health promotional benefits of JSHT.

Nevertheless, it is still noteworthy that our understanding of JSHT use has numerous limitations. The first gap is that research on the chemical composition of JSHT is currently lacking. Although several researchers have studied the individual ingredients of JSHT and identified numerous phytochemical compounds, structural modifications of the phytochemical compounds after combining all the ingredients may need to be taken into consideration. Moreover, the majority of research done on JSHT's ingredients focused on crude extracts and their abundant bioactive secondary metabolites; however, the results of these investigations are still ambiguous and inadequate. Thus, additional systematic studies on JSHT are required to identify the particular phytochemical compounds in the extracts and determine their pharmacological effects with greater precision. Further, the current health-related information on JSHT is insufficient, and its clinical value has not been sufficiently explored. Therefore, a deep and systematic phytochemical investigation of JSHT and its pharmacological properties, especially the mechanism of action, to illustrate its ethnomedicinal use and support further efficacy and safety should undoubtedly be the focus of further research.

Data availability statement

All data generated or analyzed during this study are included in this published article.

Financial support and sponsorship

Nil.

Conflicts of interest

Dr. Shinn-Zong Lin, an editorial board member at *Tzu Chi Medical Journal*, had no role in the peer review process of or decision to publish this article. The other authors declared no conflicts of interest in writing this paper.

REFERENCES

- Sarada K, Puthiyedath R, Philip A, Ravindran GC, Pavithran K. Prevalence of the use of traditional complementary and alternative medicine amongst cancer patients in a tertiary care center in Kerala, India. *J Ayurveda Integr Med* 2021;12:359-64.
- Badakhsh M, Dastras M, Sarchahi Z, Doostkami M, Mir A, Bouya S. Complementary and alternative medicine therapies and COVID-19: A systematic review. *Rev Environ Health* 2021;36:443-50.
- Zhao X, Tan X, Shi H, Xia D. Nutrition and traditional Chinese medicine (TCM): A system's theoretical perspective. *Eur J Clin Nutr* 2021;75:267-73.
- Balekundri A, Mannur V. Quality control of the traditional herbs and herbal products: A review. *Futur J Pharm Sci* 2020;6:1-9.
- Hsieh PC, Chao YC, Tsai KW, Li CH, Tzeng IS, Wu YK, et al. Efficacy and safety of complementary therapy with Jing Si Herbal tea in patients with mild-to-moderate COVID-19: A prospective cohort study. *Front Nutr* 2022;9:832321.
- Lin YJ, Chiang CY, Shibu MA, Su SH, Dass KT, Lin PY, et al. Novel Herbal Formulation Jing Si Exhibits Multiple Functions to Inhibit Replication Activity and Subsidies Viral Load of COVID-19 Variants; 2022.
- Chu S, Wang W, Zhang N, Liu T, Li J, Chu X, et al. Protective effects of 18 β -Glycyrrhetic acid against myocardial infarction: Involvement of PI3K/Akt pathway activation and inhibiting Ca²⁺ influx via L-type Ca²⁺ channels. *Food Sci Nutr* 2021;9:6831-43.
- He YK, Yao YY, Chang YN. Characterization of anthocyanins in *Perilla frutescens* var. *Acuta* extract by advanced UPLC-ESI-IT-TOF-MSn method and their anticancer bioactivity. *Molecules* 2015;20:9155-69.
- Gyawali S, Luintel S, Thapa S, Kunwar RM, Bussmann RW, Paniagua-Zambrana NY. *Anisomeles indica* (L.) Kuntze *Lamiaceae*. In: *Ethnobot Himalayas*. Springer, Cham; 2020, p 1-4.
- Cheng FJ, Huynh TK, Ko HY, Hsueh CS, Yeh YL, Wu YC, et al. *Artemisia argyi* Potentially Prevents the Infections with SARS-CoV-2 Variants; 2023.
- Xiao W, Zhou P, Wang X, Zhao R, Wang Y. Comparative characterization and immunomodulatory activities of polysaccharides extracted from the radix of *Platycodon grandiflorum* with different extraction methods. *Molecules* 2022;27:4759.
- Misra P, Datta SK, Chakrabarty D. Mutation in flower colour and shape of *Chrysanthemum morifolium* induced by γ -radiation. *Biol Plant* 2003;47:153-6.
- Romagna IS, Junges E, Ramos RF, Antonioli ZI, Bellé C. Meloidogyne incognita parasitizing *Ophiopogon japonicus* in Rio Grande do Sul state, Brazil. *Australas Plant Dis Notes* 2019;14:29.
- Liccardi F, Boscutti F, Sigura M, Tordoni E, Carpanelli A, Valecic M, et al. First report of naturalization of *Houttuynia cordata* Thunb. 1783 (*Saururaceae*) in Italy. *Rendiconti Lincei. Scienze Fisiche e Naturali* 2021;32:287-93.
- Anilkumar D, Joshi H, Nishteswar K. Review of *Glycyrrhiza glabra* (Yastimadhu)-a broad spectrum herbal drug. *Pharma Sci Monit* 2012;3:3171-95.
- Nasrin S, Islam MN, Tayab MA, Nasrin MS, Siddique MA, Emran TB, et al. Chemical profiles and pharmacological insights of *Anisomeles indica* Kuntze: An experimental chemico-biological interaction. *Biomed Pharmacother* 2022;149:112842.
- Song X, Wen X, He J, Zhao H, Li S, Wang M. Phytochemical components and biological activities of *Artemisia argyi*. *J Funct Foods* 2019;52:648-62.
- Ryu J, Nam B, Kim BR, Kim SH, Jo YD, Ahn JW, et al. Comparative analysis of phytochemical composition of gamma-irradiated mutant cultivars of *Chrysanthemum morifolium*. *Molecules* 2019;24:3003.
- Pastorino G, Cornara L, Soares S, Rodrigues F, Oliveira MB. Licorice (*Glycyrrhiza glabra*): A phytochemical and pharmacological review. *Phytother Res* 2018;32:2323-39.
- Wu Z, Deng X, Hu Q, Xiao X, Jiang J, Ma X, et al. *Houttuynia cordata* Thunb: An ethnopharmacological review. *Front Pharmacol* 2021;12:714694.
- Chen MH, Chen XJ, Wang M, Lin LG, Wang YT. *Ophiopogon japonicus* – A phytochemical, ethnomedicinal and pharmacological review. *J Ethnopharmacol* 2016;181:193-213.
- Ahmed HM. Ethnomedicinal, phytochemical and pharmacological investigations of *Perilla frutescens* (L.) Britt. *Molecules* 2018;24:102.
- Zhang L, Wang Y, Yang D, Zhang C, Zhang N, Li M, et al. *Platycodon grandiflorus* – An ethnopharmacological, phytochemical and pharmacological review. *J Ethnopharmacol* 2015;164:147-61.
- Lu PH, Tseng CW, Lee JL, Lee EY, Lin YP, Lin IH, et al. Jing Si herbal drink as a prospective adjunctive therapy for COVID-19 treatment: Molecular evidence and mechanisms. *Pharmacol Res Chinese Med* 2022;2:100024.
- Shibu MA, Lin YJ, Chiang CY, Lu CY, Goswami D, Sundhar N, et al. Novel anti-aging herbal formulation Jing Si displays pleiotropic effects against aging associated disorders. *Biomed Pharmacother* 2022;146:112427.
- Taleghani A, Emami SA, Tayarani-Najaran Z. *Artemisia*: A promising plant for the treatment of cancer. *Bioorg Med Chem* 2020;28:115180.
- Wu LK, Agarwal S, Kuo CH, Kung YL, Day CH, Lin PY, et al. *Artemisia* Leaf Extract protects against neuron toxicity by TRPML1 activation and promoting autophagy/mitophagy clearance in both *in vitro* and *in vivo* models of MPP+/MPTP-induced Parkinson's disease. *Phytomedicine* 2022;104:154250.
- Kwon YJ, Son DH, Chung TH, Lee YJ. A review of the pharmacological efficacy and safety of licorice root from corroborative clinical trial findings. *J Med Food* 2020;23:12-20.
- Bisht D, Rashid M, Arya RK, Kumar D, Chaudhary SK, Rana VS, et al. Revisiting liquorice (*Glycyrrhiza glabra* L.) as anti-inflammatory, antivirals and immunomodulators: Potential pharmacological applications with mechanistic insight. *Phytomed Plus* 2022;2:100206.
- Li PC, Wang HS, Shibu MA, Wang J, Huang SH, Wang JH, et al. Clinical course of patients with severe SARS-CoV-2 infection co-treatment with Jin Si Herbal Tea in Eastern Taiwan: A retrospective cohort study. *J Herb Med* 2022;36:100610.
- Hsieh SC, Fang SH, Rao YK, Tzeng YM. Inhibition of pro-inflammatory mediators and tumor cell proliferation by *Anisomeles indica* extracts. *J Ethnopharmacol* 2008;118:65-70.
- Shin NR, Park SH, Ko JW, Ryu HW, Jeong SH, Kim JC, et al. *Artemisia argyi* attenuates airway inflammation in lipopolysaccharide induced acute lung injury model. *Lab Anim Res* 2017;33:209-15.
- Guerrero JA, Lozano ML, Castillo J, Benavente-García O, Vicente V, Rivera J. Flavonoids inhibit platelet function through binding to the thromboxane A2 receptor. *J Thromb Haemost* 2005;3:369-76.
- Cinatl J, Morgenstern B, Bauer G, Chandra P, Rabenau H, Doerr HW. Glycyrrhizin, an active component of liquorice roots, and replication of SARS-associated coronavirus. *Lancet* 2003;361:2045-6.
- Tsai KC, Huang YC, Liaw CC, Tsai CI, Chiou CT, Lin CJ, et al. A traditional Chinese medicine formula NRICM101 to target COVID-19 through multiple pathways: A bedside-to-bench study. *Biomed Pharmacother* 2021;133:111037.
- Kou J, Yu B, Xu Q. Inhibitory effects of ethanol extract from radix *Ophiopogon japonicus* on venous thrombosis linked with its endothelium-protective and anti-adhesive activities. *Vascul Pharmacol* 2005;43:157-63.
- Tang WF, Tsai HP, Chang YH, Chang TY, Hsieh CF, Lin CY, et al. *Perilla* (*Perilla frutescens*) leaf extract inhibits SARS-CoV-2 via direct virus inactivation. *Biomed J* 2021;44:293-303.
- Kim TY, Jeon S, Jang Y, Gotina L, Won J, Ju YH, et al. *Platycodon* D, a natural component of *Platycodon grandiflorum*, prevents both

- lysosome- and TMPRSS2-driven SARS-CoV-2 infection by hindering membrane fusion. *Exp Mol Med* 2021;53:956-72.
39. Huang HC, Lien HM, Ke HJ, Chang LL, Chen CC, Chang TM. Antioxidative characteristics of *Anisomeles indica* extract and inhibitory effect of ovatodiolide on melanogenesis. *Int J Mol Sci* 2012;13:6220-35.
 40. Han HM, Kim SJ, Kim JS, Kim BH, Lee HW, Lee YT, et al. Ameliorative effects of *Artemisia argyi* Folium extract on 2,4 dinitrochlorobenzene induced atopic dermatitis like lesions in BALB/c mice. *Mol Med Rep* 2016;14:3206-14.
 41. Rigopoulos D, Kalogeromitros D, Gregoriou S, Pacouret JM, Koch C, Fisher N, et al. Randomized placebo-controlled trial of a flavonoid-rich plant extract-based cream in the treatment of rosacea. *J Eur Acad Dermatol Venereol* 2005;19:564-8.
 42. Choi KT, Kim JH, Cho HT, Lim SS, Kwak SS, Kim YJ. Dermatologic evaluation of cosmetic formulations containing *Chrysanthemum indicum* extract. *J Cosmet Dermatol* 2016;15:162-8.
 43. Grippaudo FR, Di Russo PP. Effects of topical application of B-resorcinol and glycyrrhetic acid monotherapy and in combination with fractional CO(2) laser treatment for benign hand hyperpigmentation treatment. *J Cosmet Dermatol* 2016;15:413-9.
 44. Mapoung S, Umsumarng S, Semmarath W, Arjsri P, Srisawad K, Thippraphan P, et al. Photoprotective effects of a hyperoside-enriched fraction prepared from *Houttuynia cordata* Thunb. On ultraviolet b-induced skin aging in human fibroblasts through the MAPK signaling pathway. *Plants (Basel)* 2021;10:2628.
 45. Yu X, Gao D, Qi B, Xiao X, Zhai X, Ma CW, et al. *Ophiopogon japonicus* herbal tea ameliorates oxidative stress and extends lifespan in *Caenorhabditis elegans*. *Pharmacogn Mag* 2018;14:617.
 46. Mungmai L, Preedalikit W, Pintha K, Tantipaiboonwong P, Aunsri N. Collagenase and melanogenesis inhibitory effects of *Perilla frutescens* pomace extract and its efficacy in topical cosmetic formulations. *Cosmetics* 2020;7:69.
 47. Mungmai L, Preedalikit W, Aunsri N, Amornlerdpison D. Efficacy of cosmetic formulation containing *Perilla frutescens* leaves extract for irritation and aging skin. *Biomed Pharmacol J* 2020;13:779-87.
 48. Park SJ, Lee HA, Kim JW, Lee BS, Kim EJ. *Platycodon grandiflorus* alleviates DNCB-induced atopy-like dermatitis in NC/Nga mice. *Indian J Pharmacol* 2012;44:469-74.
 49. Chiang CY, Lin YJ, Weng WT, Lin HD, Lu CY, Chen WJ, et al. Recuperative herbal formula Jing Si maintains vasculature permeability balance, regulates inflammation and assuages concomitants of "Long-COVID". *Biomed Pharmacother* 2023;163:114752.
 50. Lien HM, Huang SH, Chang CH, Huang CL, Chen CC, Chyau CC. Innovative purification method of ovatodiolide from *Anisomeles indica* to induce apoptosis in human gastric cancer cells. *Molecules* 2022;27:587.
 51. Lin CS, Bamodu OA, Kuo KT, Huang CM, Liu SC, Wang CH, et al. Investigation of ovatodiolide, a macrocyclic diterpenoid, as a potential inhibitor of oral cancer stem-like cells properties via the inhibition of the JAK2/STAT3/JARID1B signal circuit. *Phytomedicine* 2018;46:93-103.
 52. Tseng CP, Huang YL, Chang YW, Liao HR, Chen YL, Hsieh PW. Polysaccharide-containing fraction from *Artemisia argyi* inhibits tumor cell-induced platelet aggregation by blocking interaction of podoplanin with C-type lectin-like receptor 2. *J Food Drug Anal* 2020;28:115-23.
 53. Hodaie M, Rahimmalek M, Behbahani M. Anticancer drug discovery from Iranian *Chrysanthemum* cultivars through system pharmacology exploration and experimental validation. *Sci Rep* 2021;11:11767.
 54. Wang S, Shen Y, Qiu R, Chen Z, Chen Z, Chen W. 18 β -glycyrrhetic acid exhibits potent antitumor effects against colorectal cancer via inhibition of cell proliferation and migration. *Int J Oncol* 2017;51:615-24.
 55. Song X, Yin S, Zhang E, Fan L, Ye M, Zhang Y, et al. Glycycomarin exerts anti-liver cancer activity by directly targeting T-LAK cell-originated protein kinase. *Oncotarget* 2016;7:65732-43.
 56. Deng QP, Wang MJ, Zeng X, Chen GG, Huang RY. Effects of glycyrrhizin in a mouse model of lung adenocarcinoma. *Cell Physiol Biochem* 2017;41:1383-92.
 57. Subhawa S, Chewonarin T, Banjerdpongchai R. The effects of *Houttuynia cordata* Thunb and piper ribesoides wall extracts on breast carcinoma cell proliferation, migration, invasion and apoptosis. *Molecules* 2020;25:1196.
 58. Kim JM, Hwang IH, Jang IS, Kim M, Bang IS, Park SJ, et al. *Houttuynia cordata* Thunb promotes activation of HIF-1A-FOXO3 and MEF2A pathways to induce apoptosis in human HepG2 hepatocellular carcinoma cells. *Integr Cancer Ther* 2017;16:360-72.
 59. Zhang W, Zhang Q, Jiang Y, Li F, Xin H. Effects of ophiopogonin B on the proliferation and apoptosis of SGC 7901 human gastric cancer cells. *Mol Med Rep* 2016;13:4981-6.
 60. Zang QQ, Zhang L, Gao N, Huang C. Ophiopogonin D inhibits cell proliferation, causes cell cycle arrest at G2/M, and induces apoptosis in human breast carcinoma MCF-7 cells. *J Integr Med* 2016;14:51-9.
 61. Abd El-Hafeez AA, Fujimura T, Kamei R, Hirakawa N, Baba K, Ono K, et al. Synergistic tumor suppression by a *Perilla frutescens*-derived methoxyflavone and anti-cancer tyrosine kinase inhibitors in A549 human lung adenocarcinoma. *Cytotechnology* 2018;70:913-9.
 62. Zhang Y, Zhang Y, Liang J, Kuang HX, Xia YG. Exploring the effects of different processing techniques on the composition and biological activity of *Platycodon grandiflorus* (Jacq.) A.DC. by metabolomics and pharmacologic design. *J Ethnopharmacol* 2022;289:114991.
 63. Jeon D, Kim SW, Kim HS. Platycodin D, a bioactive component of *Platycodon grandiflorum*, induces cancer cell death associated with extreme vacuolation. *Anim Cells Syst (Seoul)* 2019;23:118-27.
 64. Zeggwagh NA, Farid O, Michel JB, Eddouks M. Cardiovascular effect of *Artemisia herba Alba* aqueous extract in spontaneously hypertensive rats. *Methods Find Exp Clin Pharmacol* 2008;30:375-81.
 65. Gao T, Zhu ZY, Zhou X, Xie ML. *Chrysanthemum morifolium* extract improves hypertension-induced cardiac hypertrophy in rats by reduction of blood pressure and inhibition of myocardial hypoxia inducible factor-1 α expression. *Pharm Biol* 2016;54:2895-900.
 66. Zhang L, Yang Y, Yu L, Wang Y, Liu L, Fan X. Cardioprotective effects of *Glycyrrhiza uralensis* extract against doxorubicin-induced toxicity. *Int J Toxicol* 2011;30:181-9.
 67. Jiao Y, Wang S, Jiang L, Sun X, Li J, Liu X, et al. 2-undecanone protects against fine particles-induced heart inflammation via modulating Nrf2/HO-1 and NF- κ B pathways. *Environ Toxicol* 2022;37:1642-52.
 68. Qin D, Yue R, Deng P, Wang X, Zheng Z, Lv M, et al. 8-Formylophiopogonanone B antagonizes doxorubicin-induced cardiotoxicity by suppressing heme oxygenase-1-dependent myocardial inflammation and fibrosis. *Biomed Pharmacother* 2021;140:111779.
 69. Paradee N, Utama-Ang N, Uthaipibull C, Porter JB, Garbowski MW, Srichairatanakool S. Extracts of Thai *Perilla frutescens* nutlets attenuate tumour necrosis factor- α -activated generation of microparticles, ICAM-1 and IL-6 in human endothelial cells. *Biosci Rep* 2020;40:BSR20192110.
 70. Yu L, Liu H. Perillaldehyde prevents the formations of atherosclerotic plaques through recoupling endothelial nitric oxide synthase. *J Cell Biochem* 2018;119:10204-15.
 71. Lin YC, Lin YC, Kuo WW, Shen CY, Cheng YC, Lin YM, et al. Platycodin D reverses pathological cardiac hypertrophy and fibrosis in spontaneously hypertensive rats. *Am J Chin Med* 2018;46:537-49.
 72. Kotha P, Badri KR, Nagalapuram R, Allagadda R, Chippada AR. Anti-diabetic potential of the leaves of *Anisomeles malabarica* in streptozotocin Induced Diabetic Rats. *Cell Physiol Biochem* 2017;43:1689-702.
 73. Wu X, Zhuang J, Bai Z, Guo D. *In vivo* antidiabetic activity of aqueous extract of *Artemisia argyi* (Chinese mugwort) in alloxan-induced diabetic rats. *Trop J Pharm Res* 2020;19:1487-93.
 74. Yamamoto J, Tadaishi M, Yamane T, Oishi Y, Shimizu M, Kobayashi-Hattori K. Hot water extracts of edible *Chrysanthemum*

- morifolium* Ramat. Exert antidiabetic effects in obese diabetic KK-Ay mice. *Biosci Biotechnol Biochem* 2015;79:1147-54.
75. Yamashita Y, Kishida H, Nakagawa K, Yoshioka Y, Ashida H. Liquorice flavonoid oil suppresses hyperglycaemia accompanied by skeletal muscle myocellular GLUT4 recruitment to the plasma membrane in KK-A(y) mice. *Int J Food Sci Nutr* 2019;70:294-302.
 76. Kumar M, Prasad SK, Krishnamurthy S, Hemalatha S. Antihyperglycemic activity of *Houttuynia cordata* Thunb. In streptozotocin-induced diabetic rats. *Adv Pharmacol Sci* 2014;2014:809438.
 77. Qiao Y, Jiao H, Wang F, Niu H. Ophiopogonin D of *Ophiopogon japonicus* ameliorates renal function by suppressing oxidative stress and inflammatory response in streptozotocin-induced diabetic nephropathy rats. *Braz J Med Biol Res* 2020;53:e9628.
 78. Kim DH, Kim SJ, Yu KY, Jeong SI, Kim SY. Anti-hyperglycemic effects and signaling mechanism of *Perilla frutescens* sprout extract. *Nutr Res Pract* 2018;12:20-8.
 79. Zheng J, He J, Ji B, Li Y, Zhang X. Antihyperglycemic effects of *Platycodon grandiflorum* (Jacq.) A. DC. Extract on streptozotocin-induced diabetic mice. *Plant Foods Hum Nutr* 2007;62:7-11.
 80. Lim DW, Kim YT, Jang YJ, Kim YE, Han D. Anti-obesity effect of *Artemisia capillaris* extracts in high-fat diet-induced obese rats. *Molecules* 2013;18:9241-52.
 81. Lee Y, Lee J, Lee MS, Chang E, Kim Y. *Chrysanthemum morifolium* flower extract ameliorates obesity-induced inflammation and increases the muscle mitochondria content and AMPK/SIRT1 activities in obese rats. *Nutrients* 2021;13:3660.
 82. Mirtaheri E, Namazi N, Alizadeh M, Sargheini N, Karimi S. Effects of dried licorice extract with low-calorie diet on lipid profile and atherogenic indices in overweight and obese subjects: A randomized controlled clinical trial. *Eur J Integr Med* 2015;7:287-93.
 83. Tunkamnerdthai O, Auvichayapat P, Boonmars T, Tavichakorntrakool R, Leelayuwat N. *Houttuynia cordata* Thunb. Extract attenuates dyslipidemia in high-fat-diet-induced obese rats. *Srinagarind Med J* 2013;28:199.
 84. Wang Y, Zhu Y, Ruan K, Wei H, Feng Y. MDG-1, a polysaccharide from *Ophiopogon japonicus*, prevents high fat diet-induced obesity and increases energy expenditure in mice. *Carbohydr Polym* 2014;114:183-9.
 85. Thomas SS, Kim M, Lee SJ, Cha YS. Antiobesity effects of purple perilla (*Perilla frutescens* var. *Acuta*) on adipocyte differentiation and mice fed a high-fat diet. *J Food Sci* 2018;83:2384-93.
 86. Ke W, Bonilla-Rosso G, Engel P, Wang P, Chen F, Hu X. Suppression of high-fat diet-induced obesity by *Platycodon grandiflorus* in mice is linked to changes in the gut microbiota. *J Nutr* 2020;150:2364-74.
 87. Rani PS, Rao R. Evaluation of anti-diabetic and hepatoprotective activities of aqueous and ethanolic extracts of *Anisomeles malabarica* Roots. *IJPSR* 2021;12:5079-86.
 88. Amat N, Upur H, Blazeković B. *In vivo* hepatoprotective activity of the aqueous extract of *Artemisia absinthium* L. Against chemically and immunologically induced liver injuries in mice. *J Ethnopharmacol* 2010;131:478-84.
 89. Zhou Y, Wang C, Kou J, Wang M, Rong X, Pu X, et al. Chrysanthemum flos extract alleviated acetaminophen-induced rat liver injury via inhibiting oxidative stress and apoptosis based on network pharmacology analysis. *Pharm Biol* 2021;59:1378-87.
 90. Gou SH, He M, Li BB, Zhu NY, Ni JM. Hepatoprotective effect of total flavonoids from *Glycyrrhiza uralensis* Fisch in liver injury mice. *Nat Prod Res* 2021;35:6083-7.
 91. Kang H, Koppula S. Hepatoprotective effect of *Houttuynia cordata* thunb extract against carbon tetrachloride-induced hepatic damage in mice. *Indian J Pharm Sci* 2014;76:267-73.
 92. Chen X, Tang J, Xie W, Wang J, Jin J, Ren J, et al. Protective effect of the polysaccharide from *Ophiopogon japonicus* on streptozotocin-induced diabetic rats. *Carbohydr Polym* 2013;94:378-85.
 93. Yang SY, Hong CO, Lee GP, Kim CT, Lee KW. The hepatoprotection of caffeic acid and rosmarinic acid, major compounds of *Perilla frutescens*, against t-BHP-induced oxidative liver damage. *Food Chem Toxicol* 2013;55:92-9.
 94. Ke W, Wang P, Wang X, Zhou X, Hu X, Chen F. Dietary *Platycodon grandiflorus* attenuates hepatic insulin resistance and oxidative stress in high-fat-diet induced non-alcoholic fatty liver disease. *Nutrients* 2020;12:480.
 95. Shin JM, Son YJ, Ha IJ, Erdenebileg S, Jung DS, Song DG, et al. *Artemisia argyi* extract alleviates inflammation in a DSS-induced colitis mouse model and enhances immunomodulatory effects in lymphoid tissues. *BMC Complement Med Ther* 2022;22:64.
 96. Zeng M, Li M, Chen Y, Zhang J, Cao Y, Zhang B, et al. A new bisepoxyignan dendranlignan A isolated from *Chrysanthemum flower* inhibits the production of inflammatory mediators via the TLR4 pathway in LPS-induced H9c2 cardiomyocytes. *Arch Biochem Biophys* 2020;690:108506.
 97. Yuan H, Ji WS, Wu KX, Jiao JX, Sun LH, Feng YT. Anti-inflammatory effect of diammonium glycyrrhizinate in a rat model of ulcerative colitis. *World J Gastroenterol* 2006;12:4578-81.
 98. Woranam K, Senawong G, Utaiwat S, Yunchalard S, Sattayasai J, Senawong T. Anti-inflammatory activity of the dietary supplement *Houttuynia cordata* fermentation product in RAW264.7 cells and Wistar rats. *PLoS One* 2020;15:e0230645.
 99. Zhao JW, Chen DS, Deng CS, Wang Q, Zhu W, Lin L. Evaluation of anti-inflammatory activity of compounds isolated from the rhizome of *Ophiopogon japonicus*. *BMC Complement Altern Med* 2017;17:7.
 100. Park DD, Yum HW, Zhong X, Kim SH, Kim SH, Kim DH, et al. *Perilla frutescens* extracts protects against dextran sulfate sodium-induced murine colitis: NF- κ B, STAT3, and Nrf2 as putative targets. *Front Pharmacol* 2017;8:482.
 101. Kim TY, Yoon E, Lee D, Imm JY. Antioxidant and anti-inflammatory activities of *Platycodon grandiflorum* seeds extract. *CyTA J Food* 2020;18:435-44.
 102. Kim KO, Lee D, Hiep NT, Song JH, Lee HJ, Lee D, et al. protective effect of phenolic compounds isolated from Mugwort (*Artemisia argyi*) against contrast-induced apoptosis in kidney epithelium cell line LLC-PK1. *Molecules* 2019;24:195.
 103. Xia DZ, Lv GY, Yu XF, Wang HM, Yang Q. Antagonism of total flavonoids from *Chrysanthemum morifolium* against lead induced oxidative injury in mice. *Zhongguo Zhong Yao Za Zhi* 2008;33:2803-8.
 104. Nassan MA, Soliman MM, Aldhahrani A, Althobaiti F, Alkhedaide AQ. Ameliorative impacts of *Glycyrrhiza glabra* Root extract against nephrotoxicity induced by gentamicin in mice. *Food Sci Nutr* 2021;9:3405-13.
 105. Kang C, Lee H, Hah DY, Heo JH, Kim CH, Kim E, et al. Protective Effects of *Houttuynia cordata* Thunb. On gentamicin-induced oxidative stress and nephrotoxicity in rats. *Toxicol Res* 2013;29:61-7.
 106. Sheng X, Yang Y, Liu J, Yu J, Guo Q, Guan W, et al. Ophiopogonin A alleviates hemorrhagic shock-induced renal injury via induction of Nrf2 Expression. *Front Physiol* 2020;11:619740.
 107. Kim HR, Kim SY. *Perilla frutescens* sprout extract protect renal mesangial cell dysfunction against high glucose by modulating AMPK and NADPH oxidase signaling. *Nutrients* 2019;11:356.
 108. Zhang W, Hou J, Yan X, Leng J, Li R, Zhang J, et al. *Platycodon grandiflorum* saponins ameliorate cisplatin-induced acute nephrotoxicity through the NF- κ B-mediated inflammation and PI3K/Akt/apoptosis signaling pathways. *Nutrients* 2018;10:1328.
 109. Uddin M, Abdullah-AI-Mamun M, Biswas K, Asaduzzaman M, Rahman M. Assessment of anticholinesterase activities and antioxidant potentials of *Anisomeles indica* relevant to the treatment of Alzheimer's disease. *Orient Pharm Exp Med* 2016;16:113-21.

110. Hu C, Zhang S, Chen Q, Wang R. Ovatodiolide protects ischemia-reperfusion-induced neuronal injury via microglial neuroinflammation via mediating SIRT1/NF- κ B pathway. *Brain Res Bull* 2022;180:97-107.
111. Kang JY, Park SK, Guo TJ, Ha JS, Lee DS, Kim JM, et al. Reversal of trimethyltin-induced learning and memory deficits by 3,5-dicaffeoylquinic acid. *Oxid Med Cell Longev* 2016;2016:6981595.
112. Rashidi R, Ghorbani A, Rakhshandeh H, Mousavi SH. Protective effect of *Artemisia absinthium* on 6-hydroxydopamine-induced toxicity in SH-SY5Y cell line. *Avicenna J Phytomed* 2021;11:238-46.
113. Sapkota A, Gaire BP, Cho KS, Jeon SJ, Kwon OW, Jang DS, et al. Eupatilin exerts neuroprotective effects in mice with transient focal cerebral ischemia by reducing microglial activation. *PLoS One* 2017;12:e0171479.
114. Wang H, Wang H, Cheng H, Che Z. Ameliorating effect of luteolin on memory impairment in an Alzheimer's disease model. *Mol Med Rep* 2016;13:4215-20.
115. Dong LM, WRCY, Yin TP, Li XB, Du LF, Chen CY, et al. Anti-Parkinsonian activity of *Chrysanthemum morifolium* ramat extractin mice. *Acta Nutr Sin* 2017;39:294-8.
116. Zhu Z, Qian S, Lu X, Xu C, Wang Y, Zhang X, et al. Protective properties of the extract of *Chrysanthemum* on patients with ischemic stroke. *J Healthc Eng* 2021;2021:3637456.
117. Ban JY, Park HK, Kim SK. Effect of glycyrrhizic acid on scopolamine-induced cognitive impairment in mice. *Int Neurourol J* 2020;24:S48-55.
118. Petramfar P, Hajari F, Yousefi G, Azadi S, Hamed A. Efficacy of oral administration of licorice as an adjunct therapy on improving the symptoms of patients with Parkinson's disease, A randomized double blinded clinical trial. *J Ethnopharmacol* 2020;247:112226.
119. Ravanfar P, Namazi G, Atigh M, Zafarmand S, Hamed A, Salehi A, et al. Efficacy of whole extract of licorice in neurological improvement of patients after acute ischemic stroke. *J Herb Med* 2016;6:12-7.
120. Huh E, Kim HG, Park H, Kang MS, Lee B, Oh MS. *Houttuynia cordata* improves cognitive deficits in cholinergic dysfunction Alzheimer's disease-like models. *Biomol Ther (Seoul)* 2014;22:176-83.
121. Moon JY, Kim ES, Choi SJ, Kim JI, Choi NS, Lee K, et al. Protective effect of the ethyl acetate-fraction of methanol extract of *Ophiopogon japonicus* on amyloid beta peptide-induced cytotoxicity in PC12 cells. *J Life Sci* 2019;29:173-80.
122. Lin M, Sun W, Gong W, Zhou Z, Ding Y, Hou Q. Methylophiopogonanone a protects against cerebral ischemia/reperfusion injury and attenuates blood-brain barrier disruption *in vitro*. *PLoS One* 2015;10:e0124558.
123. Cho E, Lee J, Sin JS, Kim SK, Kim CJ, Park MH, et al. Effects of *Perilla frutescens* var. *Acuta* in amyloid β toxicity and Alzheimer's disease-like pathology in 5XFAD mice. *Food Chem Toxicol* 2022;161:112847.
124. Kang HB, Kim SH, Uhm SH, Kim DK, Lee NS, Jeong YG, et al. *Perilla frutescens* leaf extract attenuates vascular dementia-associated memory deficits, neuronal damages, and microglial activation. *Curr Issues Mol Biol* 2022;44:257-72.
125. Nam Y, Shin SJ, Park YH, Kim MJ, Jeon SG, Lee H, et al. *Platycodon grandiflorum* root protects against A β -induced cognitive dysfunction and pathology in female models of Alzheimer's disease. *Antioxidants (Basel)* 2021;10:207.
126. Kim JI, Jeon SG, Kim KA, Kim JJ, Song EJ, Jeon Y, et al. Platycodon grandiflorum root extract improves learning and memory by enhancing synaptogenesis in mice hippocampus. *Nutrients* 2017;9:794.
127. Basappa G, Kumar V, Sarojini BK, Poornima DV, Gajula H, Sannabommaji T, et al. Toxicity evaluation of *Anisomeles indica* Kuntze leaf flavonoid fraction. *J Pharmacogn Nat Prod* 2016;2:1000122-32.
128. Ogbale EA, Ogundeko T, Asalu AF, Builders M, Aguiyi J. Acute toxicity studies of locally cultivated *Artemisia annua* leaf extract in Rats. *World J Pharm Sci* 2014;1864-70.
129. Li L, Gu L, Chen Z, Wang R, Ye J, Jiang H. Toxicity study of ethanolic extract of *Chrysanthemum morifolium* in rats. *J Food Sci* 2010;75:T105-9.
130. Nazari S, Rameshrad M, Hosseinzadeh H. Toxicological effects of *Glycyrrhiza glabra* (Licorice): A review. *Phytother Res* 2017;31:1635-50.
131. Zhang M, Liu X, Song J, Hu Y, Wang X, Li B. Effects of radix *Ophiopogonis* decoction on embryo-fetal development in rats. *Zhongguo Zhong Yao Za Zhi* 2010;35:2334-7.
132. Lee H, Kang R, Kim YS, Chung SI, Yoon Y. Platycodin D inhibits adipogenesis of 3T3-L1 cells by modulating Kruppel-like factor 2 and peroxisome proliferator-activated receptor gamma. *Phytother Res* 2010;24 Suppl 2:S161-7.